

UC-NRLF



\$B 262 043




4
 A COMMONPLACE-BOOK of THOUGHTS, MEMORIES, and FANCIES, Original and Selected. PART I. Ethics and Illustrative. PART II. Literature and Art. *Second Edition*, revised and corrected, with Etchings and Wood Engravings. Crown 8vo. 12s.

5
 TWO LECTURES on the SOCIAL EMPLOYMENTS of WOMEN.—*Sisters of Charity and the Communion of Labour*. New Edition, with a *Postscript* Letter on the Present Condition and Requirements of our Women of England. Fcp. 8vo. 2s.

6
 SKETCHES in CANADA, and RAMBLES among the RED MEN. 20mo. pp. 2s. 6d. cloth 97. 10 Two Parts, 1s. each.
 "THE latter work was composed as possibly a record of his own social views and
 THE TIMES, March 1888"

London: LONGMAN, GREEN, and CO. Paternoster Row.









Digitized by the Internet Archive
in 2007 with funding from
Microsoft Corporation

NAUTICAL ASTRONOMY

LONDON

PRINTED BY SPOTTISWOODE AND CO.

NEW-STREET SQUARE

THE
PROJECTION AND CALCULATION
OF
THE SPHERE

FOR YOUNG SEA OFFICERS

BEING

A COMPLETE INITIATION INTO NAUTICAL ASTRONOMY

BY S. M. SAXBY, R.N.

PRINCIPAL INSTRUCTOR OF NAVAL ENGINEERS, HER MAJESTY'S STEAM RESERVE
LATE OF CAIUS COLLEGE, CAMBRIDGE



LONDON

LONGMAN, GREEN, LONGMAN, AND ROBERTS

1861

...

For reference only

...

...

...

...

...

...

PREFACE.

SPHERIC TRIGONOMETRY is so little understood by those who are not professionally mathematicians, while the subject is so important to a maritime nation, that the question whether there be not defect in the present mode of educating nautical men demands earnest attention.

For more than a century the method of teaching what is called the "Art of Navigation" has been gradually changing, and the knowledge of *principles* has in England been consequently decreasing, while other nations, possessed of a navy and commercial marine vastly inferior to those of England, have maintained a standard which actually originated in the labours and experience of our own early nautical teachers.

It is remarkable that, while in the year 1700 no one who was ignorant of Spherics, and the principles of Nautical Astronomy, was considered a competent navigator, in the year 1860, the "epitomes" of navigation used both in the Royal Navy and the Merchant Service of this great country, not only make no pretension to the teaching of Spherics, either by calculation or construction, but one of

them (which on all other naval subjects is elaborate and accurate) even repudiates the propriety of mixing theory with practice, while certain other works insist on theory as the only medium by which to obtain a correct practical knowledge of the subject.

The true system of teaching lies between these extremes ; and this little work is an attempt to facilitate the study by sufficiently explaining general principles while adapting them to immediate practice.

A few recent works could be referred to which have already, by an advance in the right direction, whetted the mental appetite for really digestible and nourishing sustenance ; but the young naval officer and the young aspiring merchant commander of the present generation have absolutely (beyond these) no available means of improving their acquaintance with what they feel to be indisputably essential to their self-respect as navigators, inasmuch as works in publication either disregard theory altogether as unimportant to a beginner, or bewilder the student with a phalanx of formulæ, which in their very aspect so commonly suggest a series of difficulties insurmountable, that relapse into indifference is the natural consequence.

And yet naval instructors are expected to impart a knowledge of Nautical Astronomy adequate to the needs of a sailor. Such is undoubtedly done to the very limit of their powers, but few besides nautical teachers can appreciate the labour of the work of elucidating a subject like Nautical Astronomy, in the absence of well-digested plans of instruction in a printed form, in which everything would be explained *so clearly as to encourage the student, and lead him step by step up the ladder of*

knowledge, when the ascent is to be attempted away from professional assistance.

The subject of Spherics is one of such profound research and extent, that it seems to have been expected that a certain conciseness of language and solemnity of tone would best become the author who would enter publicly upon its teaching. But *familiar* description, divested of "hard words," has been so successfully attempted by an excellent mathematician of our own times while treating of sister branches of science, that the example of such men as the Rev. Harvey Goodwin (whose illustrations of general mathematics have so much benefited the Cambridge student) may safely be imitated; certainly by one, it is presumed, whose experience as a teacher of adults extends over a period of some three and thirty years.

There may be about 100,000 navigators in Great Britain, very likely there are considerably more, inasmuch as in the merchant service alone there are about 45,000 vessels afloat. It is scarcely, however, to be expected that the "wary mariner," trained as he has been to vigilance, and ever alive as he consequently is to necessity for precautions in his sea voyage, will not apply the same to his voyage of life. *It is contrary to his professional habits* to venture heedlessly among the rocks and shoals which may beset a coast to him unknown; and in like manner such are his precautions with regard to any *science* to him unknown; but give him even a tolerably clear *outline* "chart," and general "sailing directions," or even furnish him with a "hand lead" wherewith he may *feel the bottom if he cannot see it*, and there are in his occupations days and hours of thoughtful leisure in which he may be tempted to vary

his "cruise," and at least examine for himself the creeks and lagoons of science. At present these are obscured from his vision by characteristic prejudices and distrust. To remove these is then the object of this little book. Whatever may be the beauties of a science so captivating as Nautical Astronomy, the very approaches have been made toilsome from the *decay of the landmarks* which once guided the young seaman on his path. And when a straggling, casual, and mere wayfaring adventurer has accidentally gained a "peep" within the barriers, what has he seen but a frowning array of sines, secants, tangents, &c., twisted into every imaginable equational and fractional form, and distorted into a thousand labyrinthine channels, leading into deeps to him unfathomable. It is in vain to say such is a befitting initiation for a young aspirant in Nautical Science. It appals him; it drives him back to his ordinary level, discourages from other attempts to advance himself in the scale of proficiency, and throws him upon the dangerous and debilitating "consolation" that among his own class he can still "pass muster." Many indefatigable navigators *do* however brave the difficulties which present themselves, and attain a very fair footing: *the object of this book is to vastly increase their number.*

It may be asked, could one in a thousand even illustrate geometrically or by "projection" the simple question of latitude from a meridian altitude, or answer it by the use of scale and dividers? The writer would be very sorry to be compelled to publish, even from his individual experience, details of positive danger which this state of things has entailed upon the commercial world; but one thing is certain, viz. the importance of the interests of

maritime commerce is involved in the *safe transit* of its vast treasures ; such interests are paramount, and demand our best efforts in their support, — a sufficient apology for this attempt.

It is not enough that naval officers have peculiar advantages in nautical training, or that certain inducements are proffered by the Board of Trade to men capable of acquiring, or willing to contend for, “extra certificates” in the merchant service ; for not even with the latter is a knowledge of Spherics expected beyond its one simple application to great circle-sailing, and *this without any requirements whatever as to calculation*. Hence we may safely affirm that the subject of teaching Spherics seems to demand complete revision.

Nor is it sufficient that the nautical tutors of the present day perform their duties in a manner which has already loosened, as it were, the stability of an erroneous system of teaching, and which would in time restore that system to its previous sturdy and useful basis. But it is desirable to shorten such time of changeful probation, and an endeavour, in order to be successful, must be bold and radical. Navigation in these days is not as formerly required for the slow hulls of the beginning of the last century, which averaged from four to six knots an hour, but for steamers, swiftly flying against wind and tide at the rate of — (it is scarcely prudent to say *what!*). A more *ready system* of navigation is therefore called for ; and if years since it were important to the venerable structures to which allusion is made, that *they* should be navigated in the best possible method, by so much the more is it necessary that in this improved age of “clippers,”

all the efficacy of a *sound knowledge of principles* should be called into operation.

If Spherics *as a basis* of all oversea navigation was considered and found to be indispensable to the safe conduct of ships in 1680, by so much the more must a knowledge of it be essential to shipmasters in 1860.

In tracing the cause of the deterioration which the system of nautical training has undergone during the last century and a half, we shall find it shown in the preface to the still admirable and elaborate "Practice of Navigation," written by the lamented and unrequited Lieut. Raper, R.N., wherein he states that the theory and the practice are kept "*purposely distinct*;" and he adds "it is the custom generally to teach the theory first; the impression forced on me is on the contrary, that the practice is itself the best foundation for sound and rapid advancement in the theory. For he who has acquired the practice knows the nature and extent of the subject, and in proceeding to the theory he has a distinct perception of the object to be attained."

The author of this little work concurs fully in the views of Raper, wherein he advocates the all importance of practical knowledge to the student in theory; but the exclusion of theory is merely the lesser evil, and has led to the "rule of thumb" system, to the final exclusion of elucidation of principles from our best existing works on navigation on the one hand, or has engrossed it too exclusively on the other, in works intended for young sea officers; its inevitable consequence is therefore the state of things which it is the object of the writer of this to combat.

The mind of a seafaring man is so much occupied with responsibilities of duty, or as regards passengers, the crew, &c., that he has in general too little leisure for deep studies, and when on shore his necessary and reasonable repose admits of little interruption; therefore to benefit the whole class a work is wanted which, in familiar language, and in a proper blending of theory with practice can lead to a solid and respectable acquirement. The more easy the steps of knowledge can be rendered to so peculiarly situated a class as sea officers, the faster progress will he make who aims at a higher intellectual level; and, indeed, a very *gently inclined plane of science is more necessary for the navigator than for any other class of individuals.*

Such, then, this book is intended to be; such, indeed, as a young seaman may take up with a firm conviction that all essentials are fully explained; and with an interest in believing that if it be read even with the attention usually paid to a work of fiction, *as a mere pastime*, sound and valuable information will infallibly take root in his mind.

The oft-quoted assertion attributed to Euclid in his reply to one of the Ptolemies, viz. "that there is no royal road to learning," must no longer obtain among us. Not every seaman aspires to become an Archimedes, and if he did, an Alexandrian school might not be the only one and the best in which to form him.

What Sir John Herschel has done for astronomy, Faraday for chemistry, Sedgwick for geology, Arnold for classics, Hutton and Colenso for arithmetic and algebra, and Goodwin, Snowball, &c., for the study of mechanics,

may surely be done for the navigator. Hence the object of this book.

Ill understood, isolated and forbidding formulæ thrust imperiously upon an already burdened memory are a load and an oppression, while pleasant, easy illustration of principles is an enticement, and wins upon the mind until its powers are secured by the silken bonds of a willing captivity.

The British Government are seconding the community in their struggles for a better *middle class* education. Shall then those whose perilous avocations demand especial intelligence, and to whose hands and *heads* we trust our lives and properties, and above all our national honour and the defence of our hearths, shall *these alone* be excluded from the social arena of progress, and as mere spectators see the palm of proficiency in the grasp of a foreigner?

S. M. S.

CONTENTS.

	Page
PROJECTION : —	
Orthographic	2
Mercator's	7
Stereographic	8
GENERAL REMARKS ON TRIGONOMETRY	10
Definition of an angle	15
Geometrical theorems	16
Ratios	21
Natural sines	22
LOGARITHMS, the nature of	23
Computation of	29
NAUTICAL ASTRONOMY	31
Lines of the sphere	32
Spheric triangle	34
General diagram of daily astronomical phenomena	36
SPHERIC PROJECTION	38
Plane scale	39
PROB. I. To make an angle so that the angular point shall be at the centre of the primitive	39
II. To lay off any number of degrees on a right circle	40
III. To draw an oblique circle through any point lying within the primitive circle	40
IV. To draw an oblique circle through any two points	41
V. To draw a small circle parallel to the primitive at any distance from it	41
VI. To draw a parallel circle at a given distance from a right circle, or at any distance about a given point at the primitive	42
VII. To find the pole of an oblique circle	43

	Page
SPHERIC PROJECTION (<i>continued</i>):—	
PROB. VIII. To draw a great circle through any given point so as to make any desired angle at the primitive	43
IX. To draw a small circle through a given point which shall be at a given distance from a right circle	44
X. To draw an oblique circle perpendicular to a given oblique circle	45
XI. To draw a great circle which shall make any angle with the primitive	45
XII. To measure any part of an oblique circle	46
XIII. To measure an angle at the primitive	47
XIV. To measure an angle which is not at the primitive	47
CONSTRUCTION OF SPHERIC TRIANGLES	
To find the latitude of a place	51
To find the apparent time	56
To find an azimuth	58
CALCULATION OF SPHERIC TRIANGLES	
Right angled	60
The five circular parts	62
Quadrantal spheric triangles	69
Oblique spheric triangles	69
CASE 1. Two sides and an opposite angle	71
2. Two angles and an opposite side	74
3. Two sides and an included angle	76
4. Two angles and an included side	80
5. Three sides	83
6. Three angles	85
NAUTICAL ASTRONOMY:—	
Ex. 1, 2, 3, 4, 5. To find latitude	87
6, 7, 8. To find apparent time	96
9, 10. To find an azimuth	100
11. To find a great circle course	102
12. To find a great circle distance	104
13. To find the latitude of a great circle vertex	105
14. To find an altitude	107
15. To find an amplitude	108
16. To find the time of daybreak	109
17. To find the time of rising of any celestial body	111

Erratum.

Page 24, line 12, *for* $1 + 1 + \frac{1}{1.2}$ *read* $1 + 1 + \frac{1}{1.2}$



TRIGONOMETRY

AND

NAUTICAL ASTRONOMY.

WHILE plane trigonometry in its application to the practice of navigation, is so well set forth in the usual epitomes and other nautical works, that little need be said herein upon the subject, beyond a few remarks on certain of its fundamental principles, no one can properly comprehend even the very elementary principles of nautical astronomy, without a better insight into spherical trigonometry than is to be obtained from any work at present accessible to the sea officer.

PROJECTION.

The study of spherics is too often undertaken in ignorance of a method of constructing a spheric angle by scale. Such methods (for there are several) are called "projections," a general term in spherics, signifying the transferring of spaces from rounded surfaces to flat or "plane" ones, in the forms in which the eye would trace them, according to their assumed relative position.

Distortion must of necessity accompany the projection

of a spheric surface upon a plane, and all methods used in accomplishing this are not equally good.


1. For example, the gnomonic projection presents peculiar advantages in dialling, and also in great circle or tangent sailing; while the orthographic projection is used by the astronomer in the delineation of eclipses, the transits of heavenly bodies, &c. Mercator's is exceedingly useful in the construction of sea charts; while the stereographic is more applicable than all others to the purposes of nautical astronomy, in consequence of all its parts being either arcs of circles or straight lines. The globular is used by map engravers, the scenographic for perspective, and need not occupy the attention of the nautical astronomer.

2. Our consideration, then, will be restricted to the orthographic, Mercator's and stereographic projections; merely first noting that in the gnomonic projection, the eye is supposed to be at the centre of the sphere, viewing the meridians as straight lines; and in this projection the shortest distance on the globe between two places is represented by the shortest distance between the two corresponding points on the flat surface.

ORTHOGRAPHIC.

3. In the orthographic projection, we, for the occasion, must suppose the eye to be placed at an immense distance,


Fig. 1.



and as viewing a sphere as if it were a mere disc, such as the sun or moon appears to be, the visual rays, as 1, 2, 3, &c., in Fig. 1, being supposed to be parallel to each other; in such case parallels of latitude in a right sphere would appear (as *c, d, x*, in Fig. 2) to be straight lines, and separated in the proportion of the divisions on what we call the line of sines upon the scale; while the meridians lying near the diameter would appear to be at nearly their actual distances asunder, but would be crowded as they lie nearer to the periphery as at *a* (Fig. 1).*

4. Supposing, further, that a globe of immense size had the usual lines of the sphere marked upon it, it would, as

Fig. 2.



seen by the eye at (if possible) an extreme distance, have the appearance of Fig. 2, in which all the meridians are ellipses and *not* arcs of circles; this would readily be understood by holding a small toy globe at arm's length, with the polar diameter in a vertical direction.

5. There is, perhaps, at the present time, no work in pub-

* Many illustrations might be given from one crowded diagram; but as plainness is so desirable in a work like this, it is better to give separate diagrams with the text therewith connected.

lication which would much assist the ordinary student in projecting the orthographic sphere. The methods given, and the explanations accompanying them, are beyond the comprehension of a beginner; and are, moreover, so sufficiently troublesome to the practised draughtsman as to cause him in most cases merely to mark off the orthographic distances of the meridians on *the equator only*, and draw them as arcs of circles. This may even be seen in the diagrams illustrating the works of our greatest philosophers. Hence orthographic plates will seldom bear the inspection of the mathematician. It is not therefore extraordinary that the orthographic projection has been so little used for nautical purposes, until the writer suggested its introduction (as it will be further explained), to relieve the stereographic projection of certain disabilities in its application to the purposes of navigation.


6. The following directions for constructing an orthographic hemisphere, in what is called a "right sphere" (having the poles in the *circumference* as distinguished from a "parallel sphere" in which the poles are at the *centre*, and from an "oblique sphere" in which the poles are *neither* at the circumference nor at the centre) are given in detail, the method not having been yet published, and being the one long used by the writer. Its utility may at once be inferred from the circumstance of its requiring neither tangents nor secants, but simply the line of sines and scale of chords. It should be borne in mind that in Fig. 2 the parallels all appear as straight lines, like the equator; and if the line, *a b*, in its divisions forms the line of sines as seen at the diameter of Fig. 1, any corresponding portions of the other parallels must do the same, because they are supposed to be viewed from the same point; hence, *c d*, and *e f*, Fig. 2, &c., are each crossed by meridians, at distances proportional to the natural sines of their respective latitudes. In other words, d^2 would

be a semi-minor axis of the meridian $\approx 2n$, or the cosine of its meridian. Every nautical astronomer, therefore, will do well to have a scale prepared for general use, similar to the one at Fig. 3. It is formed by making the lines BC and BA meet at any angle (the nearer 90° the more convenient), and laying off on BA the distances B 15, B 30, &c., to B 90, from any scale of sines; then join C 15, C 30, &c., and *parallel* to BA draw any number of lines through the figure, and the scale is complete.

Fig. 2 may be drawn from this scale in the following manner:—

Take AB from the scale (Fig. 3), as a radius, and describe a circle, drawing two diameters right-angled at the centre. Take a straight edge of paper and lay it at BA, Fig. 3; transfer on to it the distances, B 15, B 30, &c., up


Fig. 3.



to 90; lay these off on Fig. 2, as $b_1, b_2, \&c.$, up to ba , and also from b towards z , from b to y , and from b to n . Through bz and bn draw parallels as in Fig. 2; then take the length of any *half* parallels, as cd , on a straight edge of paper, find its distance on the scale (Fig. 3), as at cd , and having copied the divisions (as was done at BA in dividing ab , Fig. 2), lay off these on Fig. 2, as $d_1, d_2, d_3, \&c.$: proceed thus with each half parallel, and points will be accurately obtained through which the whole of the desired meridians may be constructed.

7. If it be required to project the sphere on the plane of the equator, or the “parallel sphere,” proceed as follows. From the scale of sines (Fig. 3) take BA as radius, and with it draw a circle $abcd$, quartering it, as before, at right angles, as in Fig. 4. If it be desirable to draw meridians through every 10° , or every 15° (suppose the latter) take the division of the line of sines, Fig. 3 (as before), apply them from e towards b , in Fig. 4; and with the centre e , and distances $e1$, $e2$, &c., describe circles; then lay off from a to b , b to c , c to d , and d to a , every 15th degree from a scale of chords corresponding

Fig. 4.



to the radius (or by dividing either quadrant into six equal parts); join these several points with the centre, and these form the meridians, the poles of the primitive circle being at the centre.

8. This mode of construction is particularly useful in projecting places in the polar regions, or places of heavenly bodies when near the poles of the ecliptic or equator; in such cases the least distortions being at and near the centre of the figure; but is most especially valuable in general questions of nautical astronomy, where the data

lie near the prime vertical, such as altitudes, &c., at hours which are near 6 A.M. or 6 P.M., while for other periods of the day or night nearer to noon or midnight, the stereographic projection has its advantages.

MERCATOR'S.

9. Mercator's projection is too well known by nautical men to require much mention here, but its properties may be thus very briefly stated. On a globe it will be noticed that the actual measured length of degrees of longitude diminishes as they recede from the equator towards either pole.

In about the year 1590, a Mr. Wright, of Caius College, Cambridge, conceived the notion that he could on paper conveniently compensate this contraction of the degrees of longitude (in placing the round upon a flat surface), by *expanding* each degree of latitude in proportion to its corresponding degree of longitude in that latitude; by this means he projected the meridians and parallels as straight lines; hence, in this all rhumbs (or compass bearings) cross the meridians at equal angles, and a ship's course is laid down as a straight line. But this projection greatly distorts the outline and figures of places lying far from the equator. The miles in a degree of latitude are to the miles in a degree of longitude as radius is to cosine of latitude in which the degree of longitude is situated. For example:—Required the length in measured nautical miles of a degree of longitude in latitude 50° .

In Fig. 5, making hypotenuse radius, we have (by plane trigonometry)

$$\text{rad AC} : 60 :: \cos 50^\circ : \text{AB} = 38' \cdot 57 \text{ miles of long;}$$

or in Fig. 6, making base radius

$\sec 50^\circ : 60 :: \text{rad } A B : 38' \cdot 57 \text{ miles of long.}$

Fig. 5.





Fig. 6



STEREOGRAPHIC.

10. The stereographic projection (which every navigator will feel the greatest pleasure and advantage in comprehending if he read the following with ordinary attention), enters largely into the daily work and interest of every one who has command of a ship.


In this projection the eye is supposed to be at some part of the earth's surface, say, upon the equator, and it would in such case see the meridians of the further hemisphere, as they would appear if produced on to a flat surface tangential to the opposite point of the equator which would be exactly under the eye. The following (Fig. 7) will best illustrate this; in which we suppose the globe to be transparent, and resting on a point of its equator upon a table; the eye being at *a*, and viewing the meridians, *c 5*, *c 4*, &c., as they would appear to it if their intersection of the equator, *a b D*, were produced to the surface of the table.

GENERAL REMARKS ON TRIGONOMETRY.

11. A few elementary hints, which, although essential to the proper understanding of the subject, are not always sufficiently explained to a beginner, and thereby materially retard his progress, will properly precede the consideration of spheric construction.

The following (Fig. 8) is called the trigonometrical "canon" (a word merely signifying a collection of mathematical truths or formulæ), and from it are derived all the terms and rules used and practised in trigonometry.

Fig. 8.



In the above,

AB is called the radius of the circle (of course AK, AI, AH, and AE, are also radii. Euc. I. def. 15.)

BC is called a tangent (always of the opposite angle A), and is so called from Latin, *tangere* to touch, because it only *touches* but does not cut the circle.

AC is called a secant [always of the angle between it

and radius, therefore of \angle (angle) A], so called from Latin, *secare* to cut, because it cuts through the circle.

DE is called a sine, because it lies in the *hollow*, or *bosom*, of the curve EBL (Latin, *sinus*).

FE is called a cosine, or sine of an \angle which is complementary to another, or required to make up 90° . Thus DE is the sine of the arc EB, and FE is the sine of arc EH, which is the complement of EB (for $HE + EB = 90^\circ$). Therefore FE is called the cosine of EB, and is equal to AD, because FE and AD are drawn parallel, and ED is perpendicular to both.

GH is, in like manner, the tangent of HE, or cotangent of EB.


AG is, in like manner, the secant of HE, or cosecant of EB.

HI is called a line of chords, from its serving, as it were, to tie or confine the ends of the arc HnI .

DB is called a versed sine, and is the "height of the segment" EBLDE.

12. Every circle is supposed to be divided into 360 degrees (marked 360°). If, therefore, in the following figure, (9) PZ equals 90° or a quadrant, it is plain that if Dx also equals 90° , the word "degree" refers to no measure of length, but merely signifies the 360th part of a circle, whatever the size of that circle may be; and, therefore, a degree may be of *any* length. As, however, degrees enter into calculations, *some definite value* of them *must* evidently be necessary; and, consequently, geometers express the *value* of degrees by taking any two lines from those given in the trigo-

Fig. 9.




nometrical canon (Fig. 8), and consider the *length* of one *as compared* with the *length* of another in the same triangle: so that we use the terms sine, tangent, secant, &c., as referred generally to the radius of the circle, considering the length of radius to be 1 or 10, &c. (inches, feet, miles, leagues, &c., *at will*): this will be further illustrated.

13. It has long been customary to call a *line*, as ED (Fig. 8), a sine, or as BC, a tangent; but such is only correct when the length of a radius of a circle is known or understood. It is generally useful to describe the sine, &c., as fractions, thus, $\frac{ED}{AE}$ (Fig. 8), as they express fairly the

value referred to. Anticipating by a few pages the question of proportion, it may here be noted that a vulgar fraction is a "ratio" or proportion in itself, and is deduced from a triangle. Thus, when we speak of $\frac{3}{4}$ ths of anything we refer to some magnitude which can only be appreciated by considering the fraction $\frac{3}{4}$ in relation to its integer or whole number "one." As this whole number itself, expressed as a fraction, is $\frac{4}{4}$ ths, $\frac{5}{5}$ ths, $\frac{6}{6}$ ths, &c.; and when, therefore, we speak of $\frac{3}{4}$ ths, we express a "ratio," meaning as 3 is to 4 (or symbolically 3 : 4), so then we express the value of

Fig. 10.



degrees by using ratios, and comparing them with the radius of the circle, which it is convenient to do by a number capable of decimal division, for obvious reasons (such as 1, 10, 100, &c.); and as at least one side of a plane triangle is always given, we are at liberty to compare this with the length of what is thus called the sine, tangent, secant, &c., of an angle, and

hence the length of the arc itself.

14. For further example, in Fig. 10, let BD be what is

commonly called the *sine* of the angle A. We describe its value by saying it is as the perpendicular is to radius, and write it thus: $\frac{\text{perp}}{\text{rad}}$ or from the figure $\frac{BD}{CB}$.

In like manner the other fundamental trigonometrical ratios are represented by fractions thus:—

$$\begin{aligned} \frac{\text{perp}}{\text{rad}} &= \frac{PN}{CN} \text{ is an expression for the tangent of } \angle A \\ \frac{\text{hyp}}{\text{rad}} &= \frac{CP}{CN} \quad \text{,,} \quad \text{secant of } \angle A \\ \frac{\text{rad}}{\text{perp}} &= \frac{CN}{PN} \quad \text{,,} \quad \text{cotang } \angle A \\ \frac{\text{hyp}}{\text{rad}} &= \frac{CP}{PN} \quad \text{,,} \quad \text{cosec } \angle A \\ \frac{\text{rad}}{\text{hyp}} &= \frac{CN}{CP} \quad \text{,,} \quad \text{cosine } \angle A \end{aligned}$$

From the above it will be seen that certain ratios are *reciprocals*; for instance:—

$$\begin{aligned} \text{sine} &= \frac{PN}{CP} \quad \text{and cosec} = \frac{CP}{PN} \\ \text{tang} &= \frac{PN}{CN} \quad \text{and cotang} = \frac{CN}{PN} \\ \text{sec} &= \frac{CP}{CN} \quad \text{and cosine} = \frac{CN}{CP} \end{aligned}$$

15. Therefore, in works on logarithms, when we want the secant of an angle we can find it by subtracting its cosine from 20 (the diameter of a circle whose radius is 10), and to find the log sine we subtract the log cosec from 20; and to find the log tang we subtract the log cotang from 20, &c. Other useful deductions may be made, such as to find the log tang: add 10 to the log sine, and from the

sum subtract the log cos (or log tang = $\frac{\log \text{ sine} + 10}{\log \text{ cos}}$), and to find the log cotang add 10 to the log cos, and from the sum subtract the log sine (or log cotang = $\frac{\log \text{ cos} + 10}{\log \text{ sine}}$), &c.; so that the values of the six fundamental ratios may be expressed thus:—

$$\begin{array}{ll} \text{sine} & = \frac{1}{\text{cosec}} & \text{cosine} & = \frac{1}{\text{sec}} \\ \text{tang} & = \frac{1}{\text{cot}} & \text{cot} & = \frac{1}{\text{tang}} \\ \text{sec} & = \frac{1}{\text{cosine}} & \text{cosec} & = \frac{1}{\text{sine}} \end{array}$$

N.B.—The unit here meaning 1 diameter = 2 radii each of 10, or diameter = 20.

16. This, however, which forms the elementary base of a proper knowledge of trigonometry, is not *absolutely essential* to the navigator, whose practical operations in plane trigonometry may be performed in total ignorance of principles, by dint of mere intelligence and skill from repetition; but in like manner does the blacksmith strike with the face of the hammer, and not with the handle, and would not probably perform his work more effectually if he were, previous to every blow, to calculate the force required to fashion his heated iron. But this must be remembered: a mathematical “blacksmith” would probably give *fewer blows*, because he would better know how to make each stroke tell, *from bringing the face of the hammer to bear on the iron in the best direction* with the greatest effect. In like manner the mathematical navigator will obtain his result in the shortest method.

It is beneath the dignity of a British sea officer to be content with mere knowledge of the use of formulæ. After reading this little book he may be safely advised to take up

“Jeans’s Trigonometry,” or some such *small work* on the subject.


DEFINITION OF AN ANGLE.

17. Before proceeding it may be well to explain what is really meant by an angle: that such explanation is necessary cannot be denied. A work of this description, which is designed as a mere stepping-stone to study, must needs adopt assertions without proofs, for fear of alarming the timid who *desire* improvement, but who yet doubt their own powers. It may, however, be safely asserted, that since our proofs are deduced mainly from the Books of Euclid a knowledge of his system of proving should be imparted at the earliest opportunity. In works upon navigation generally more extracts from Euclid are given than the sea officer thinks necessary for his satisfactory working, and too few to satisfy his after-desire of research; while the Books of Euclid themselves are supposed to be too heavy an undertaking for any but a schoolboy having no other employment than study. These are delusions. A groundwork in mathematics well laid is a continual source of mental profit and *amusement*. There is no limit (but the powers of mortal intellects) to the structure which may be raised upon it. A very long acquaintance with the subject of teaching can only lead to a belief that whenever mathematical study is to any mind found to be repulsive, it may be suspected that the individual student *has not had its details sufficiently explained*. Thousands upon thousands can work an equation by logarithms who have but an indistinct notion of what is really meant by an angle; and it may cheer the student when he sees that we may go even to the “dreaded” Euclid to obtain a full and clear comprehension even of this trifle.

For instance, he says among his definitions (Book I. def.

8): "A plane angle is the inclination of two lines to one another in a plane, which meet together, but are not in the same direction." And (Book I. def. 9), "A plane rectilinear angle is the inclination of two right (or straight) lines to one another, which meet together, but are not in the same right line;" so that, in the following figure the right

Fig. 11.



line AB meets the right line AC at the point A, and the "angle" is the inclination of these two lines as measured in degrees upon *any* circle drawn from A as a centre cutting these two lines. For instance, *de*, or *fg*, or *hi*, is each the measure of the "angle A" in degrees, 20 degrees meaning $\frac{20}{360}$ of *any* circle drawn round the point A.


GEOMETRICAL THEOREMS.

18. A few of Euclid's theorems may here be introduced with advantage.

Book I. XIII.—The angles which one right line makes with another upon one side of it, are either two right angles, or are together equal to two right angles. Departing from the precise language of absolute and complete proof (for obvious reasons), we may say that the semicircle HEB contains 180 degrees. If EG be perpendicular to AB, the arcs HE and EB being equal will each contain 90°, but BC is less than 90°, being, say, 55°; then EC must be 35°, and EH being 90°, CH will be 125°. Now, EC is


called the *complement* of CB, and HC is called the *supplement* of CB.

Fig. 12.



19. Book I. XV. tells us that if two right lines cut one another the vertical or opposite angles shall be equal. Thus, the angles CEA and BED are equal to each other, as are also CEB and AED; the angle CEA means the angle at the point E formed by the two lines CE EA. (We always put the letter *indicating the point* between the others.) Now, from the XIIIth Proposition, Book I., as above, it is evident that the two contiguous angles HGC and CGB equal 180 degrees; so in Fig. 13 $ac + cb$ equal 180 degrees, and $bd + da$ form the other 180 degrees.


Fig. 13.



20. Euc. I. Prop. XXIX.—If a right line falls upon two parallel right lines, it makes the alternate angles equal to one another, &c. &c.; so that, indeed, AGE, CHE, FHD,

FGB , are equal to each other, being in this case about 28 degrees, while EGB , EHD , FHC , and FGA are also equal, being about 152° ,—the circles render this apparent.

Fig. 14.




To satisfy this it needs only that AB and CD be precisely parallel.

21. Euc. I. Prop. XIX. — The greater angle of every triangle is subtended by the greater side, or has the greater side opposite to it.

The arcs of the circles drawn about each of the angular

Fig. 15.




points with an equal radius show at once that, for instance, the small angle B 30° is opposite to the smaller side AC , &c.

22. Euc. VI. Prop. VIII. — In a right-angled triangle if a perpendicular be drawn from the right angle to the base, the triangles on each side of it are similar to the whole triangle and to one another (that is, triangle ADC , triangle BDC , and triangle ACB are similar).

Now, by *similar* triangles we mean triangles which have

the three angles in the one equal to three angles respectively in the other; and although their opposite sides may be of different lengths, they are nevertheless proportional: thus, by the figure, if we lay off the distance CD at Bc , and draw ac parallel to CD , we shall find the triangle Bca equal in its angles to CDA , and its sides proportional to triangle BCD , that is, Bc will be to ac as BD is to DC ($Bc : ac :: BD : DC$), &c. &c., and here (by I., XXIX.), because BC falls across the two parallel lines ac , CD , the angles Bac and BCD are equal. (And this is the

Fig. 16.




way in which one proposition of Euclid rests for proof upon others which precede. As an example, proof of the above proposition could only be made with *mathematical accuracy* by reference to 34 propositions of Book I., 10 of Book V., and 3 of Book VI.; in all 47 *propositions*, besides *definitions, axioms, and postulates*, repetitions, &c.).

23. Euc. III. Prop. XX.—The angle at the centre of a circle is double the angle at the circumference upon the same base, that is, upon the same part of the circumference.

In the triangle ADC , the side AD equals DC , therefore, as equal sides are opposite to equal angles (as deduced from Book I. Prop. XVIII.), the angle DAC equals DCA . But Prop. XXXII. Book I. implies that the angle EDA equals DAC and DCA together; let ACB and


$A D B$ be two angles standing upon the same base $A B$; therefore, as $A D E$ is the double of $A C D$, and by like reasoning $E D B$ would be the double of $E C B$, so must the whole angle $A D B$ be the double of the angle $A C B$.

Fig. 17.



24. Euc. I. Prop. XXXII.—If any side of a triangle be produced, the exterior angle is equal to the two interior and opposite angles; and the three interior angles of any triangle are equal to two right angles.

Fig. 18.



Reference to Prop. XXIX. Book I. will show that the angle CAB will equal the angle DBE , because the right line CB falls on the two parallels AC and BD ; also, that the angle ACB will equal CBD . Therefore the angle CBE , which is exterior to the triangle, and is made up of the two angles DBE and DBC , is equal to the two interior and *opposite* angles of the triangle (the angle ABC being the *adjacent* angle). And it is also evident that the two “opposite” angles, *together with* the “adjacent” angle form the *interior* angles of the triangle) are equal to the

exterior and adjacent angles together, therefore (by I. XIII.) are equal to 180° or two right angles.

The above will be sufficient to give a general notion of the importance of Euclid.

25. Every triangle has three sides, and in lettering a right-angled triangle it is customary to place B at the right angle and to make AC designate the hypotenuse.

RATIOS.

26. The rules of Plane Trigonometry do not fall within the present compass of this work; they are to be found in all works on navigation. It may, however, be remarked that the rules given therein for the working of questions in navigation, take for instance—

as diff lat
is to rad
so is dep
to tang course,

are in the form of a proportion or ratio (proportion is the equality of ratios)—three things, as in what is called the Rule of Three, being given to find a fourth. Every conceivable arithmetical calculation is the working of a proportion, or the comparison of ratios. When we say 5 times 8 make 40, we mean to say that $1 : 5 :: 8 : 40$, or, fractionally, $\frac{1}{5}, \frac{8}{40}$; and these fractions form what is called an “equation,” for $\frac{1}{5} = \frac{8}{40}$. But Euclid, Book VI. 16, demonstrates that, if four straight lines (or quantities) be proportionals, the product of the means (or middle terms) of such proportion shall be equal to the product of the extremes (or first and last terms), as from the above $1 \times 40 = 5 \times 8$; and, as another example, if 3 coils of rope cost 20 shillings, 6 coils will cost 40, or $3 : 20 :: 6 : 40$; that is $3 \times 40 = 20 \times 6$.

27. Euclid, further, in Book V. (Def. 13 to 17), shows that such proportions may be varied by division, conversion, inversion, alternation, &c., so that the last example admits of being varied in form; for


$$\begin{aligned} \text{As } 20 : 40 :: 3 : 6 \\ 6 : 40 :: 3 : 20 \\ 40 : 6 :: 20 : 3, \text{ \&c.} \end{aligned}$$

So far, therefore, as the calculation of straight lines is concerned (which may be put to represent by their proportionate *lengths* any proportionate *quantities*), the work of calculation becomes easy; but in entering upon the calculation of angles, it should be shown that common arithmetic fails altogether in its powers to readily solve all the parts of a trigonometrical figure. Nor is the difficulty properly explained in elementary works of the present day.

NATURAL SINES.

28. If we examine Fig. 19, in which the circumference is divided into equal parts of ten degrees each, we shall see in the line CD a scale of what are called "Natural Sines." C30 being equal to a 30, &c. The divisions on

Fig. 19.



CD, moreover, are not equal; for C 10 is much larger than the distance 40 to 50; and, indeed, the sine of 30° is exactly in length half the sine of 90° , while the natural

number 30° is only one third of 90° ; and we shall also find that the secant of 60° is equal to twice the sine of 90° , while the natural number 60 is two thirds of 90; that the tangent of 45° is equal to the sine of 90° , &c.

LOGARITHMS.

29. To obviate the above inconvenience, an *artificial* system of numbers was sought for by mathematicians; and, happily, Lord Napier, about 250 years since, gave the world his System of Logarithms; and our astonishment is excited when we learn that this great discovery was made nearly half a century before what has since been called the "Logarithmic Series" was invented.

NATURE OF LOGARITHMS.

30. The very word *Logarithm* is a stumbling-block to many; but it is easy to show that, although the groundwork is so very little known to many who use logarithms, a few hints put in familiar language will not merely gratify a laudable curiosity, but pleasantly assist in further investigation.

31. Lord Napier based his system of logarithms upon the following infinite series, in which it will be seen that values of fractions are systematically diminished by adding increasing multipliers to the denominators.

$$1 + \frac{1}{1} + \frac{1}{1 \cdot 2} + \frac{1}{1 \cdot 2 \cdot 3} + \frac{1}{1 \cdot 2 \cdot 3 \cdot 4} + \frac{1}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5} + \&c.$$

Any one who understands vulgar fractions can resolve these into the following:—

$$1 + 1 + \frac{1}{2} + \frac{1}{6} + \frac{1}{24} + \frac{1}{120} + \&c.$$

Of course the series might be extended to great length.

and increased accuracy, but the above is enough for our purpose. Now, if we reduce the above fractions to a common denominator, we get

$$2 + \frac{60}{120} + \frac{20}{120} + \frac{5}{120} + \frac{1}{120}, \text{ \&c.};$$

and adding numerators we get $2\frac{86}{120}$, which, reduced to decimals, becomes = 2.71666, an *approximate* base of the Napierian logarithms, which, if extended to further terms, and the division be made as usually shown in works on logarithms, becomes (as under) 2.7182818, &c., which, so far as it extends, is the *true* base of the Napierian logarithmic system; thus:—

$$\begin{array}{rcll} 1 + 1 + 1 \cdot 2 & = & 2\frac{1}{2} & = 2 \cdot 5 \\ \frac{1}{1 \cdot 2 \cdot 3} & = & \frac{1}{6} & = \cdot 166666666 \\ \frac{1}{1 \cdot 2 \cdot 3 \cdot 4} & = & \frac{1}{24} & = \cdot 041666666 \\ \frac{1}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5} & = & \frac{1}{120} & = \cdot 008333333 \\ \frac{1}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6} & = & \frac{1}{720} & = \cdot 001388888 \\ \frac{1}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 7} & = & \frac{1}{5040} & = \cdot 0001984126 \\ \frac{1}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 7 \cdot 8} & = & \frac{1}{40320} & = \cdot 0000248015 \\ \frac{1}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 7 \cdot 8 \cdot 9} & = & \frac{1}{362880} & = \cdot 0000027557 \\ \frac{1}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 7 \cdot 8 \cdot 9 \cdot 10} & = & \frac{1}{3628800} & = \cdot 0000002755 \\ & & & \hline & & & 2 \cdot 7182818 \quad \&c. \end{array}$$

Now, the Napierian system, arising, as above shown, from

so simple an infinite series, and one which is so easily remembered, is made the basis of all other systems.

32. Any number may be taken as a base; let us at random take the number 3. Then, as logarithms are defined to be *a series of numbers in arithmetical progression, placed opposite to and corresponding with another series in geometrical progression, and so placed that 0 in the logarithmic stands opposite 1 in the geometric*—we can easily form a skeleton system based on the number 3, as under:—

Natural Numbers.	Geometrical.		Logarithms.
1 =	1	0·000000
3 ¹ =	3	1·000000
3 ² =	9 (3 × 3)	2·000000
3 ³ =	27 (3 × 3 × 3)	3·000000
3 ⁴ =	81 (3 × 3 × 3 × 3)	4·000000
3 ⁵ =	243 &c. &c.	5·000000
3 ⁶ =	729	6·000000
3 ⁷ =	2187	7·000000
3 ⁸ =	6561	8·000000
3 ⁹ =	19683	9·000000
3 ¹⁰ =	59049	10·000000

33. To prove that the above is really a table of logarithms, let us attempt calculation by it as we would by the table in common use: remembering the rules, that—

In logarithms we *multiply* numbers by *adding* their logarithms, and we *divide* numbers by *subtracting* their logarithms. Suppose, for example, we desire to multiply 729 by 81.

In the above table
 the log of 729 is 6
 log of 81 is 4
 the sum 10 = log of the answer, 59049

By arithmetic.

$$\begin{array}{r}
 729 \\
 81 \\
 \hline
 729 \\
 5832 \\
 \hline
 59049 \\
 \hline
 \end{array}$$

And again, to divide 2187 by 243,

By arithmetic.

$$\begin{array}{r}
 243)2187(9 \\
 \underline{2187} \\
 0
 \end{array}$$

By logarithms.

$$\begin{array}{l}
 \log \text{ of } 2187 \text{ is } 7 \\
 \log \text{ of } 243 \text{ is } 5 \\
 \hline
 \text{the difference } 2 = \log \text{ of } 9.
 \end{array}$$

And further, to extract the cube root of 19683. This is done by dividing the logarithm of the number by the index of the power.

$$\begin{array}{l}
 \log \text{ of } 19683 = 9, \text{ and } \frac{9}{3} \text{ the index} = 3 \text{ which is the log of } 27; \\
 \text{or, as written } \sqrt[3]{19683} = 27
 \end{array}$$

and again, to raise the number 9 to the fourth power: multiply the log by the number of the index of the power; thus—

$$\begin{array}{r}
 \log \text{ of } 9 \text{ is } 2 \\
 \text{index } 4 \\
 \hline
 8 = \log \text{ of the number } 6561; \\
 \hline
 \end{array}$$

$$\text{or } 9^4 = 6561.$$

34. A table of the above description has, however, serious defects; the greatest is apparently the want of analogy between the number of figures in the whole number and the index of the logarithm, as will be shown immediately. The index or *characteristic* of the logarithm, is the in-

teger, or whole number, and the decimal is called the *mantissa*. To facilitate calculation by logarithms, Mr. Henry Briggs, a contemporary of Lord Napier, published at Cambridge, in 1615, a system having for its base the number 10, the root of our decimal scale of notation, in which the powers of the number 10 are shown by merely adding to unity as many figures as are "indicated," by what are therefore aptly called the indices of different powers, as we see in the following skeleton table to the base 10. Here, again, the logarithm is merely the *index of the power*, while it indicates absolutely the number of figures, *less one*, in the whole number to which it corresponds. This is the common system of logarithms.

35. The above table (32) only gives the logarithms of numbers which are multiples or powers of 10, but we might, for instance, require to know the log of 270 — which would evidently lie between the log of 100 and the log of 1000; its index, however, would be 2 (because the number contains three figures), together with a decimal or mantissa, and we see in a more extended table it would be as represented by the log 2·23044.

Natural Numbers.	Logarithms.
$10^0 = 1$	0·000000 &c.
$10^1 = 10$	1·000000
$10^2 = 100$	2·000000
$10^3 = 1000$	3·000000
$10^4 = 10000$	4·000000
$10^5 = 100000$	5·000000
$10^6 = 1000000$	6·000000
$10^7 = 10000000$	7·000000
$10^8 = 100000000$	8·000000
$10^9 = 1000000000$	9·000000
$10^{10} = 10000000000$	10·000000

36. It is important to add, that if the natural number be a vulgar fraction, such as $\frac{5}{8}$, we may (because it means 5 divided by 8), subtract the log of the denominator from that of the numerator (increased by unity if necessary) thus—

$$\begin{array}{r} \log \text{ of } 5 = 0.698970 \\ \text{,, } \text{,, } 8 = 0.903090 \\ \hline 1.795880 = \text{dec. fraction } .625. \end{array} \quad \text{Proof } \begin{array}{r} 8)5000 \\ \cdot 625 \end{array}$$

It is obvious, therefore, that it would have been as simple to have reduced the vulgar fraction to its decimal at once, and then taken its logarithm.

37. But we borrowed an unit in subtracting, therefore the resulting 9 was absolutely .9 (decimal 9) or minus 1 (written $\bar{1}$). (A word here to those who are not “well up” in decimal arithmetic; be advised and lose not a day in “brushing up” a little. It is not, however, likely, that any one having sufficient interest in the subject to enable him to read this little book thus far, will do otherwise.) It will then be easily seen that it would, in the above example, have been better to borrow 10 than 1, writing the resulting index 9 as *minus* 9 (-9). Another example: multiply 100.6 by .1006.

$$\begin{array}{r} \log \text{ of } 100.6 = 2.002598 \\ \text{,, } \text{,, } .1006 = -9.002598 \\ \hline 1.005196 \end{array}$$

casting off the borrowed ten it will be 1.005196, equal to the number 10.12; this is a more simple plan than writing the log of .1006, as $\bar{1}.002598$, and operating algebraically.

The following abstract will have its uses, and illustrate the above. (The number 1006 is taken at random, any number may be substituted.)

Natural Numbers.	Logarithms.
1006	3·002598
100·6	2·002598
10·06	1·002598
1·006	0·002598
·1006	−9·002598
·01006	−8·002528
·001006	−7·002598
&c.	

N.B.—All works on logarithms have rules attached, for taking out numbers, whether representing linear or angular quantities.

COMPUTATION OF LOGARITHMS.

38. We have seen (34) that only logarithms which have a certain base are conveniently applicable to practical purposes, and that the Napierian system is the most simply obtained from a *series*, which gives its base 2·718281828, &c. This is commonly called the *natural* or hyperbolic system, and is written thus—

Log_e 2·718281828, &c., while the decimal or Brigg's or the common system is written log₁₀ (read, log to the base 10, or log to the base ε).

We use the Napierian system as a foundation of our common system in the following deduced formula:—

$$\left. \begin{array}{l} \text{The common log} \\ \text{of any number} \end{array} \right\} = \frac{\text{Nap. log of number}}{\text{the Nap. log of 10}} = \frac{\text{Nap. log. of number}}{2·3025851} =$$

$$\frac{1}{2·3025851} = ·43429448 \times \text{Nap. log of the number.}$$

Hence, to construct the common logarithm of any number, we use a number which may be called a *Napierian constant*; it is the double of the above equals ·43429448 (which is the *modulus* of the common system), and equals ·86858896.

It is here quite unnecessary to give the algebraic reasons why we use the following infinite series: it is enough for our purpose to say that in it we have a series, by which we may compute the logs of all natural numbers, and this without knowing the log of any previous number. It is this:—

$$\log P = 2M \left\{ \frac{P-1}{P+1} + \frac{1}{3} \left(\frac{P-1}{P+1} \right)^3 + \frac{1}{5} \left(\frac{P-1}{P+1} \right)^5 + \frac{1}{7} \left(\frac{P-1}{P+1} \right)^7 + \&c. \right\}$$

Now, if we let P represent the number whose log we require, say the number 2, and M the modulus, .43429448, the above will, in figures, be as under:—

$$\log 2 = 2(.43429448) \left\{ \frac{2-1}{2+1} + \frac{1}{3} \left(\frac{2-1}{2+1} \right)^3 + \frac{1}{5} \left(\frac{2-1}{2+1} \right)^5 + \frac{1}{7} \left(\frac{2-1}{2+1} \right)^7 + \&c. \right\}$$

$$\log 2 = .86858896 \left\{ \frac{1}{3} + \frac{1}{3} \left(\frac{1}{3} \right)^3 + \frac{1}{5} \left(\frac{1}{3} \right)^5 + \frac{1}{7} \left(\frac{1}{3} \right)^7 + \&c. \right\}$$

(or, as reduced)

$$\log 2 = .86858896 \left(\frac{1}{3} + \frac{1}{81} + \frac{1}{1215} + \frac{1}{15309} + \&c. \right)$$

(or decimally)

$$\log 2 = .86858896 (.333333 + .012347 + .000823 + .000065 + \&c.)$$

(or, after addition)

$$\log 2 = .86858896 \times .346568$$

(or, after multiplication)

$$\log 2 = .301025 \text{ approximately, but if the series be extended}$$

$$\log 2 = .3010300 \text{ as we find in our tables of common logarithms.}$$

NAUTICAL ASTRONOMY.

39. Before entering upon precise rules for construction and calculation of spheric angles, a few remarks upon the subject of nautical astronomy itself will lighten the labour of the student.

The general complaint of those who have “looked into” spherics, is, that although they have been taught to work spheric angles, they do not understand the principles sufficiently to be able to apply their knowledge with confidence to ordinary or rather extraordinary work.

If, say they, we could always see the figures illustrating our questions, were it only in the mind’s eye, it would assist us in obtaining solutions, and increase our interest in the study; it would also very materially help our memories.

Too generally, however, a spheric triangle is drawn by hand, without any reference whatever to its adaptation to the data of the question under consideration; hence much theoretical difficulty arises from the consideration of angles, as acute or obtuse, and of complements, supplements, &c. If in the study of spherics, the maturing of the reasoning powers is to be our main object (as in teaching Euclid for the mere logic of its reasonings), present works on the subject are abundant. But if the study is to be undertaken by those circumstanced like sea officers in general, whose object is to master enough of the principles to enable them to pursue their professional avocations in mathematics, with confidence and success, an abridged arrangement of scientific facts suitable to their purpose will doubly benefit them, inasmuch as not only will their nautical work be more accurately and more readily performed, but the having once obtained a well-grounded acquirement in principles, will render it less difficult for

them to employ their few hours of leisure, which some services permit, in pleasant advancement.

. **40.** The special object of the following explanations is then, to lead the young seaman to the most agreeable part of a navigator's study, viz., "construction." Not that in practice he will be required to actually draw his diagrams, *but a knowledge of construction will greatly aid him in calculation.*


LINES OF THE SPHERE.

41. To give a general notion of the imaginary lines of the sphere, we will suppose that I am standing on the coast, looking seaward towards the west. The north would in such case evidently be on my right hand, the south on my left hand, and the east would be behind me. I might imagine a point over my head called the zenith, and a point below my feet called the nadir. The distance of the zenith and nadir would, of course, be unlimited; but let us for the sake of precise illustration limit it to any distance, say, 1000 yards; in such case I must consider the horizon to the north and south also limited to the same distance. Now, suppose further, the meridian of the place on which I am standing to be a line drawn from the north point of the horizon, up over my head through the zenith point, and down precisely to the south point of the horizon, this would give me a semicircle. And again, imagine lines drawn from the point over my head (zenith) downwards, so as to cut each point of the compass at the horizon; these would be called azimuth circles.

Let me now further suppose that I walk backwards in a line due east, until I see the figure my imagination has been constructing with, say, a radius of 1000 yards; it

would, if visible, appear precisely as the following figure, 20 (if drawn on the stereographic projection, which alone will be used, with a slight exception, in the following demonstration of spherics), the centre marked west (W). being the point on which I had been previously standing.


Fig. 20.



Suppose, again, that the centre of this figure was situated in latitude 50° N., and for illustration, that the polar star is exactly at the north polar point of the heavens, or exactly over the north pole of the earth (which it is not by about $1\frac{1}{2}$ degrees), let me now, as in Fig. 21, divide my figure from north to Z, and from Z to south into degrees, 90 in each quadrant; and through the 50th degree from north or N. I place the pole of the heavens, being at about the spot at which I should see the polar star at night in the heavens, while standing at the centre of the figure. I would next imagine a line connecting this pole with the centre of the figure, and drawing another line at right angles to it from W to Q, the latter would represent the part of the equator above the horizon, SN, because every part of it would be 90° from the pole. It has already been shown that the great circle, NZS, is a meridian; it is also to an observer at W, the 12 o'clock "*hour circle*;" because, supposing the sun to be in north declination (or distance north of the equator), and rising at the point of the horizon marked *c*, it would, between

its rising and noon, seem to describe the small arc, cd , until, being at d , on the meridian of the place at noon, it would descend from d towards c , where it would "set" below the horizon. Thus we see that the point W answers either for east or west. In Figs. 20 and 21, the azimuths are drawn to every point of the compass, and whichever azimuth circle cuts the horizon at the point of the sun's setting would be its true bearing at such sunset; in Fig. 21, it would set at about W.N.W. But if PZS is an hour circle, so PW would be another (viz., the 6 o'clock hour circle), and we may conceive others to be drawn intermediate. Py is therefore one, and represents the hour circle of about half-past 2 P.M. or $\frac{1}{2}$ past 9 A.M., while O the intersection of dc and POy , would be the sun's place at that time; and it is, for example, the work of "spherics" to calculate the proportions of the spheric triangle ZOP , PO being evidently the "polar distance," and ZO the zenith distance, ZP the co-latitude, &c.

Fig. 21.




SPHERIC TRIANGLE.

42. We may now, by another figure, 22, show that what is meant by a spheric triangle is really a part of the *surface* of a solid globe, bounded by three arcs of great circles, as the triangle ABC . The angles being the in-

clinations of the planes of the great circle to each other, and the lengths of the sides have always reference to the angles they make at the centre of the solid figure, although only the triangle itself is usually shown in diagrams.

Fig. 22.



Having now, it is presumed, given a correct idea of what is to be understood by "spherics," it will be pleasant to see how this very interesting branch of science is illustrative of daily ordinary phenomena. If we inquire as to what extent of information might be obtained from very simple data, the reply will be highly encouraging, because we shall find that the drawing of a few curves by *very easy* rules (which are about to be fully explained further on) open to us a vast and satisfactory insight into astronomy itself.


GENERAL DIAGRAM OF DAILY ASTRONOMICAL PHENOMENA.

43. Let it be supposed that we are in latitude 50° N., that the date is the 20th May, and the time of day 10 in the morning.

N.B.—On the 20th May the sun's declination or distance from the equator would be about 20° N.

Figure 23 is drawn from these data, and gives us the following information accordingly :—

Fig. 23.



Data. - Lat 50° N. = PN
 Dec 20° N. = dE or OI
 Time 10 A.M. = LQPO

- SN would be the horizon.
- OB ,, the sun's altitude.
- OZ ,, ,, zenith distance.
- OP ,, ,, polar distance.

$E p$	would be a parallel of altitude.
Z	the zenith.
K	nadir.
N	north part of the horizon.
S	south.
C	east or west part of the horizon.
$O d$	sun's distance from the meridian.
ZOB	azimuth circle.
SB	azimuth from south.
BN	north.
CB	amplitude from C (east or west).
Zd	meridional zenith distance.
Sd	altitude at noon (meridional altitude).
PGH	six o'clock, hour circle.
POH	ten o'clock, or 2 P.M. hour circle.
ZC	prime vertical.
$d y$	declination, or apparent path in the heavens for the day.
$d A$	half the length of the day.
$A y$	night.
y	sun's place at midnight.
A	rising or setting.
d	noon.
O	10 A.M., or 2 P.M.
G	6 o'clock.
F	when on the prime vertical.
$\angle QPR$	time of the sun's rising or setting.
AT	the limit of duration of twilight.
$d O y$	parallel of declination, 20th May.
$w B x$	21st December.
Bw	half the length of the shortest winter's day.
Bx	longest winter's night.
Nx	the sun's distance below the horizon on 21st December at midnight.

$S w$	would be	the sun's meridian altitude on 21st December.
$N P$	„	the latitude of the place (or height of the pole).
$Z P$	„	complement of latitude.
$A G$	„	ascensional difference.

SPHERIC PROJECTION.

44. The general terms used in nautical astronomy being thus understood, it remains to illustrate “spheric projection;” but as we mean to explain as we go, let us advance warily. Only those problems which are absolutely essential to sea officers will be at first given. The remainder may possibly follow in a supplementary volume for the assistance of those who desire a more extensive acquaintance with the subject.

In the following figures the eye is supposed to be opposite the centre, which point is called the “pole” of the primitive or boundary circle; but the word pole will not henceforward in this book signify anything more than a point exactly 90° from some great circle.

Circles are either great or small, not so much from their dimensions as from their *position on the sphere*. None but great circles can divide a sphere into two equal parts, their planes cutting the centre. Small circles are those whose planes do not cut the centre, but divide the sphere into two unequal parts. Small circles are also called parallel circles, because their planes are parallel to the plane of the equator (of this kind are parallels of latitude, of declination, of altitude, &c.).

Before explaining the problems, the following is to be specially remembered, viz., all the measures used in spherics, such as sines, tangents, &c., are taken from parts of the circle; and in case the student may have forgotten the construction of the plane scale, its formation may be

seen in the accompanying diagram, from which all measures in the succeeding figures are drawn.

45. PLANE SCALE.





Fig. 24.

PROBLEMS.

46. PROB. I.—*To make an angle so that the angular point shall be at the centre of the primitive (say 50°).*


From centre O through 50° on the primitive draw Oa if the arc be already divided. If not divided, make the angle aOH equal 50° by the use of a scale of chords, laying 50° from H to a (first drawing the circle with a radius of 60° from the scale, as in all cases).

Fig. 25.



47. PROB. II.—*To lay off any number of degrees on a right circle (say 60°).*

Fig. 26.




1st (say upon OZ from the centre). Take 60° from the scale of semi-tangents and lay off from the centre to a .

2nd (say upon XO from X towards O). On the scale of semi-tangents count 60° backwards from 90° , and lay it off from X towards O , at b .

48. PROB. III.—*To draw an oblique circle through any point lying within the primitive circle.*

Fig. 27.




Connect the point (as at A) with the primitive H Z R X, and from the point (say H) at the primitive draw a diameter H O R; draw also another diameter at right angles to the first, as Z O X. Bisect the line A H, and produce (or lengthen) the bisecting line till it cuts O X at x ; then x will be the centre of an oblique great circle which will cut the point A.

49. PROB. IV.—*To draw an oblique circle through any two points, say through point I and point A.*

Through either point (say A) draw a diameter as H A O R. Draw Z X at right angles to it at the centre. Join A Z. Make A Z y a right angle at Z. The intersection of Z y on

Fig. 28.




H R produced will give a third point, y . Join $I y$ and $A I$, and bisect $Z y$ and $A I$, the lines of intersection will meet at x , which will be the centre of the great circle A I B.

50. PROB. V.—*To draw a small circle parallel to the primitive at any distance from it (say 40°).*

With its complement 50° from the scale of semi-tangents

and centre O, draw a circle as abc , and it will be 40° from the primitive.


Fig. 29.



51. PROB. VI.—*To draw a parallel circle at any given distance from a right circle (say at 50° from HR), or at any distance about a given point at the primitive (say at 40° from Z .)*

1st. Lay off 50° from the scale of chords from H to a ,

Fig. 30.




also from R to c , and 50° from the scale of semi-tangents from the centre to b . Then through the three points abc describe a circle as in Prob. IV. Or, take the secant of $(90 - 50^\circ) = 40^\circ$. Lay off this distance from O to d , and then, with the tangent of 40° and d as a centre, draw abc , the small circle required. Whether the point be b , or e , or f , &c., use the same means.

52. PROB. VII.—*To find the pole of an oblique circle (say of ZcX).*

Draw and produce Xc till it cuts the primitive. From a lay off the distance of 90° (or HZ) from a past Z to b .

Fig. 31.




Join bX ; and where this line cuts the radius OR (or at p) will be the pole of ZcX . Or thus: Measure CO on the scale of semitangents, and lay off its complement from O towards R, as at p , the pole.

53. PROB. VIII.—*To draw a great circle through any given point so as to make any desired angle at the primitive (say 40° and through the point d).*

With centre O and tangent 40° describe an arc as ab ;

with centre d and secant 40° describe another arc which cuts the first at x ; then x is the centre of the oblique circle ZdX , and it is drawn accordingly.

Fig. 32.




Note.—Diameters are always at right angles to the primitive.

54. PROB. IX.—*To draw a small circle through a given point (x), which shall be at a given distance from a right circle (say at 40°).*

Take the secant of 50° (the complement of the given

Fig. 33.




distance) and from centre O sweep an arc as $p q$; cross this with the tangent of 50° laid off from the point x . Then will n be the centre of the small circle $a d b$. Join $O n$, and draw the diameters $c e$ and $f g$. Then $g b$ and $f a$ will each measure 40° on the scale of chords, and $O d$ will measure 40° on the scale of semi-tangents.

55. PROB. X.—*To draw an oblique circle perpendicular to a given oblique circle.*

Find the pole p of the given circle $Z a X$ (Prob. VIII.). Draw any diameter at pleasure, say $b O c$; through $c p b$ draw

Fig. 34.




a great circle (Prob. IV.), and it shall be perpendicular to $Z a X$. If the perpendicular be required to pass through any point, as a , through the two points a and p , describe an oblique circle by Prob. IV.

56. PROB. XI.—*To draw a great circle which shall make an angle of, say 30° , with the primitive.*

Draw a diameter from Z at right angles to HR from the point Z as a centre, and with the chord of 60° from the plane scale by which the circle was drawn describe the arc $O E G$. From the centre O make $O E$ (on $O E G$) equal to 30° on

the scale of chords. Join ZE and produce to x , and x will be the centre of the oblique circle ZYX, and it measures 30° on the scale of semi-tangents, *counting from 90° on the*


Fig. 35.



scale, or from H towards O. Thus any oblique circle may be drawn by taking the number of degrees from the scale of secants; for example, the radius of the oblique circle 30° (according to the problem) is the secant of 30° .

57. PROB. XII.—*To measure any part of an oblique circle (as ab in $HabR$).*

Fig. 36.




Find the pole of the oblique circle as at p (by Prob. VII.). Join pa and produce to c , and join pb and produce to d , and the measure cd on the scale of chords will be the measure of ab .

Note.—By this problem any number of degrees may be laid on an oblique circle.

58. PROB. XIII.—*To measure an angle at the primitive, as HZa .*

N.B.—The angle at the primitive is always measured on a right circle which lies 90° distant, or which passes through the pole p of the oblique circle. Apply the distance Ha

Fig. 37.



in the dividers to the scale of semi-tangents, counting backwards from 90° . Then Ha is the measure of HZa and = 50° .


N.B.—This problem is very useful to the navigator,

59. PROB. XIV.—*To measure the angle ZaR .*

Having found the pole of ZaX (by Prob. VII.) to be at p , and the pole of HaR to be at p , join ap and produce

it to b , and join ap' and produce it to c , and the distance bc measured on the scale of chords (or 68°) will be the measure of the angle ZaR or HaX .

Fig. 38.



60. It must be remembered that the pole of a right circle is always at the primitive; thus the pole of ZX is at H or R (being at a distance of 90°), and the pole of HR is at Z or X .

N.B.—The right circle appears as a *diameter* in the stereographic projection.

CONSTRUCTION OF SPHERIC TRIANGLES.

61. In proceeding to the *construction* of spheric triangles the navigator must bear in mind that it is convenient to make arcs of latitude, declination, altitude, &c., to occupy the same relative positions in the sphere — that is to say, latitude is *always* on the primitive, &c., or in other words, he had better use a *right* sphere, as above, instead of an *oblique* sphere.

Declination is *always* a small circle parallel to the equator.

Altitude is *always* a small circle parallel to the horizon. Indeed, the lines, as they are drawn on Fig. 23, the illustrative diagram, ought to be perfectly understood and remembered.

In all the figures following the *same letters* will designate the same parts. Thus—

HR will always represent the horizon.

PS the polar axis of the earth as prolonged or *produced* to the heavens (of which the primitive is the imaginary limit, *the earth itself being now supposed to be the very small point at the centre of each figure*).

ZN the zenith and nadir.

EQ the equator.


dc a parallel of declination.

ab a parallel of altitude.

x the position of the heavenly body.

While the answers to the problems will be indicated by a *thick line* at the part of the figure where the answer is to be measured.

Fig. 39.



The line HR will consequently exactly coincide in posi-

tion with, and represent the wooden horizon of, the artificial globe (Fig. 39), while the other lines of the sphere correspond also with those on the globe, but seen as slightly distorted by the nature of the projection.

62. Those who possess a globe will do well to compare the following problems with the lines on it.

It is necessary to remember that a mere spheric triangle may be formed from any three parts given, and either a side or angle may be placed according to convenience in construction. But the limiting of certain data to certain parts of the projection is a mere conventional rule, in order to simplify the study to the minds of those who have neither time nor inclination to perfectly master the whole doctrine of spherics, but who desire a mere knowledge of its principles and practice as applicable to the wants of the navigator.

MEMORANDA.

63. All *azimuth circles* meet at the zenith and cut the horizon at right angles, and are measured along it.

All *hour circles* meet at the poles of the world, which are points on the primitive 90° from the equator.

A parallel of declination is a small circle which the sun or a heavenly body seems to describe round the pole.

The spheric figure in general use, although a hemisphere, really represents the whole sphere, inasmuch as the hour circles merely imply *time from noon*, A.M. or P.M.; and, consequently, the hour circle for 10 A.M. answers for 2 P.M., each being two hours from noon. In like manner with azimuth circles, the point next to south may be either S. by E. or S. by W. according as we consider the centre as the east or west point.

Amplitude is distance of an azimuth circle from W. or E., as measured on the horizon.

Azimuth is distance of an azimuth circle from N. or S., as measured on the horizon.

Latitude is distance from the equator.

Longitude is distance from the meridian which passes through Greenwich Observatory.

64. As the numbers used in the following constructions are merely intended to serve the purpose of illustration, answers are given to the nearest degree only.

The usually-occurring questions in nautical astronomy will first be answered by projection, and afterwards (107) the same figure will be repeated when working by calculation, such additions and arrangements being made to them as the process of calculation requires, in order to adapt them to it.


TO FIND THE LATITUDE OF A PLACE.

- 65.** I.—Given, meridian altitude sun's centre, 60° .
 „ declination 20° N.
 (Observer north of the sun.)

Draw the circle with the chord of 60° .

Draw two diameters, H R and Z N, at right angles to

Fig. 40.




same direction northward of the equator (because declination is north), where these intersect will be the centre from which with the tangent of 70° describe the small circle dc (by Prob. IX.); through this centre and the centre of the primitive draw the polar axis, PS , and the measure, RP , on the scale of chords will be the latitude.

- 67.** III.—Given, declination, 10° S.
 „ altitude . 50° .
 „ time . 10 A.M.

N.B.—In south latitude (being south of the sun) draw the primitive with the chord of 60° .

In the primitive assume a point S (above what is intended to be the south point of the horizon), and quarter the circle.

Fig. 42.



Draw SxP , two hours or 30° from the primitive (Prob. XI.).

Draw $dc = 10^\circ$ on the south side of the equator by Prob. IX. 10° from EQ .

Then with the secant from centre O , and tangent from

centre x of the co-altitude = 40° , find the centre of al (Prob. IX.), and draw it with tangent 40° as a radius; through the centre of this parallel and the centre of circle draw a diameter, ZON , and another, HOR , at right angles to it, and the distance, HS , will be the latitude, 40° south.

68. IV. — Given, altitude of a celestial body on the meridian, below the pole, say,—


Altitude of α Lyræ . . . 20° below the pole.

Declination of α Lyræ . . . $38\frac{1}{2}^\circ$ N.

(Observer south of the star.)

Let x be the star's place, the declination being $38\frac{1}{2}^\circ$, the equator $E Q$, will be $38\frac{1}{2}^\circ$ south of it, as measured on

Fig. 43.




the scale of chords, and the altitude, 20° , will give the horizon at R 20° below x . Draw HR and EQ , and diameters at right angles to each, and the height of the pole P above R will be the latitude, $71\frac{1}{2}^\circ$. The parallels of declination and altitude may be drawn by Prob. IX.

Draw the parallel of altitude, 20° , by Prob. IX., then x will be the sun's place.

Draw the parallel of declination, $d c$, through the point x , by Prob. IX.

By Prob. IV. draw $P x S$ through the three points, and the angle $Z P x$ will be the hour circle, and is measured on $E y$, a diameter at right angles to $P S$, and is equal to 45° , or three hours from noon, *westerly*, by the equator, or 3 P.M.

Fig. 46.



- 72.** VIII.—Given, latitude . 21° N.
 „ declination 20° S.
 „ altitude . 30° .

Draw $H R$ and $Z N$.

Make $R P$ equal to 21° from the scale of chords.

Draw $P S$ and $E Q$.

Draw the parallel of declination, $d c$, by Prob. IX.

Draw the parallel of altitude, $a l$, by Prob. IX., and the intersection x will be the sun's place; through the points $P x$ and S draw the oblique circle (by Prob. IV.), and the angle $Z P x$ will be the hour angle, and measured on



Draw SP and EQ .

Draw declination north by Prob. IX. (as dc), and

Draw al , the parallel of altitude, 20° by Prob. IX., then x is the sun's place.


Through the three points, Zx and N (by Prob. IV), draw an oblique circle, ZyN , and the angle yZR will be the azimuthal angle = 45° , as measured at R to y , on the semi-tangents *backwards* from 90° on the scale.

- 74. X.**—Given, latitude . 21° N.
 „ time . . 9 A.M.
 „ declination 20° S.

Draw HR and ZN .

Make RP equal to the latitude 21° .

Draw POS and EQ .



Draw the hour circle, three hours or 45° from noon (or the primitive), by Prob. XI., as Ey .

Draw the parallel of declination, dc , south (by Prob. IX.), the point of intersection, x will be the sun's place; through the points ZxN draw an oblique azimuth circle by Prob. IV., and the angle xZR will be the azimuth from north, or N. 130° E., and the angle HZy will be the azimuth from south, or S. 50° E.

CALCULATION OF SPHERIC TRIANGLES.

75. In proceeding to the calculation of spheric triangles, we notice that such are either right angled (*i.e.* having one angle equal to 90°), quadrantal (*i.e.* having one side equal to 90°), or oblique (*i.e.* having neither an angle nor a side equal to 90°).

1. RIGHT-ANGLED SPHERIC TRIANGLES.

76. Every triangle, as in plane triangles, has “six parts,” viz., three sides and three angles; and any three of these being given, the rest may be found by proportion. But in a right-angled spheric triangle two parts only need be given besides the right angle.

77. In calculating parts of a triangle, whether plane or spherical, *we shall often save much trouble if we consider, first, whether of the three things or parts given any two of them are a side and an opposite angle.* When such is the case the “rule of sines,” as it is called, founded on the fundamental theorem that “the sides of a triangle are in proportion to the sines of their opposite angles” is peculiarly simple.

To find a Side.

78. RULE.—As the sine of any given angle is to the sine of its opposite side, so is the sine of any other given angle to the sine of its opposite side.

To find an Angle.

79. RULE.—As any given side is to the sine of its opposite angle, so is any other given side to the sine of its opposite angle.


It is not, as already declared, the purpose of this book to do more than give a plain but substantially practical

introduction to the study of spherics, leaving the argumentative proofs of various theorems to the few works on the subject which are already before the public, or to a succeeding volume.

80. But that which above has been called a “fundamental” rule deserves, in passing, a little attention, be it only to encourage the student towards further research; as in this he will see the simplicity of the study of geometry when it is approached by a proper path.

In the following plane triangle ABC bisect each side and erect perpendiculars; they will meet in O, the centre

Fig. 50.



(N.B.—Lengths and angles are marked in order that the student may easily verify by logarithms).

of the circumscribing circle. Join OA, OB, and OC. Now, by reference to page 19, we shall find that Euclid, in Book III., Prob. XX., proves that “an angle at the centre of a circle is double the angle at the circumference upon the same base;” therefore, in the above figure, the angle AOB is double the angle ACB; but by construction AD is the half of AB; similarly, the angle AOD is the half of the angle AOB; therefore, the angle AOD equals the angle ACB. Now, AB, the base of angles AOB and ACB, is a chord of the arc ACB, and AD being half of AB (being by

definition called a “sine”), subtends the angle AOD or ACB.

Hence we find AD equals the sine of the angle AOD—equals the sine of the angle ACB.

By taking another base as AC, and another as CB, we shall, by the same method of demonstration, find that EC is equal to the sine of angle CBA, and also that FB is equal to sine of the angle BAC, and putting a for the side BC, and b for the side AC, and c for the side AB, we shall have the following equations:—

$$\begin{aligned}\frac{1}{2} a &= \text{sine } A \text{ (i.e. sine of } \angle A) \\ \frac{1}{2} b &= \text{sine } B \\ \frac{1}{2} c &= \text{sine } C\end{aligned}$$

And by combination:—

$$\begin{aligned}\frac{1}{2} a : \frac{1}{2} b &:: \text{sine } A : \text{sine } B, \\ \text{Or, } a : b &:: \text{sine } A : \text{sine } B, \text{ \&c.} \\ \text{Or, } a : \text{sine } A &:: b : \text{sine } B, \text{ \&c.}\end{aligned}$$

Thus the sides are in proportion to the sines of these opposite angles.

A number of useful formulæ, which seem to wear so forbidding an aspect in ordinary works upon Trigonometry, are really nothing more than easily obtained deductions from the above.

THE FIVE CIRCULAR PARTS.

81. When, however, the calculation of right-angled spheric triangles cannot fall under the rule above given (from having no angle and opposite side given), a method invented by Lord Napier and published in 1614, and which is called the “Circular Parts” (or because the right angle is never considered one of them, is called also the “Five Circular Parts”) is singularly applicable to all cases which can occur.

82. Any one of these five circular parts may be considered the *middle* part, the parts joining thereto being called *extremes conjunct*; but the parts which are *separated* by an angle or a side are called *extremes disjunct*.

N.B. The right angle does not separate its two containing sides.


83. In every case, then, as three things must enter into every consideration of proportion, viz. the two given parts (excluding the right angle) and the part required, one must be called a *middle*, while the others are considered as *conjunct* or *disjunct*, as the case may be, but in using the part in computation it must be remembered that—

“When angles or hypotenuse
Among the parts you trace,
Their complements
Or supplements
Must always take their place.”

(A quaint rhyme or two may aid the memory.)

84. The manner in which the equations are formed from which we derive the proportions may be thus explained:—

Fig. 51.



Draw any spheric triangle ABC.

Then, because the right circle AB passes through O the

pole of the primitive of which one side BC of the triangle is an arc, the angle B is a right angle.

The "Five Parts" are, therefore (83):—

1. The complement of the hypotenuse AC.
2. The side AB
3. The side CB
4. The complement of angle A
5. The complement of angle C.

Now, suppose, in the above triangle, the sides AC and AB are given to find the $\angle A$.

The middle part must be so selected as to make the other parts either disjunct or conjunct (not one conjunct and the other disjunct). The middle part in this triangle will evidently be the $\angle A$, as the two given sides include it, and because they *join it* they will be extremes conjunct.

N.B. The hypotenuse is always the side opposite the right angle. Calculation then depends on the following universal, or as it has been long called the "catholic proposition," viz. :—

85. The sine of the middle part multiplied into radius is reciprocally proportional with the tangents of extremes conjunct, and with the cosines of extremes disjunct.

Expressed as an equation it would stand thus:—

$$\text{sine of middle} \times \text{radius} = \text{tan extr conjunct} \times \text{other extr conjunct.}$$

Remembering that of four numbers in proportion, the product of the means always equals the product of the extremes (see page 21), we may vary the above as follows, viz. :—

$$\begin{aligned} \text{radius} : \text{tan extr conjunct} &:: \text{tan other extr conjunct} : \text{sine of middle.} \\ \text{or, tan extr conjunct} : \text{radius} &:: \text{side of middle} : \text{tan other extr conj.} \end{aligned}$$

If the extremes are disjunct we have as follows :—

$$\text{sine of middle} \times \text{radius} = \text{cos extr disjunct} \times \text{cos other extr disjunct.}$$

or, radius : cos extr disjunct : : cos other extr disjunct : sine of middle ;
 or, cos extr disjunct : radius : : sine of middle : cos other extr disjunct.

It follows, then, that as we can only want, in any case of right-angled spheric trigonometry, to find either a middle part or an extreme, the following four simple formulæ are all-sufficient:—

If extremes are conjunct.

RULE A. Sine of middle = $\frac{\text{tang extr conjunct} \times \text{tang other extr conjunct}}{\text{radius}}$.

RULE B. Tang extr conjunct = $\frac{\text{radius} \times \text{sine middle part}}{\text{tang other extr. conjunct}}$.

If extremes are disjunct.

RULE C. Sine of middle = $\frac{\text{cos extr disjunct} \times \text{cos other extr disjunct}}{\text{radius}}$.

RULE D. Cos. extr disjunct = $\frac{\text{radius} \times \text{middle part}}{\text{cos other extr disjunct}}$.

Or, as adapted at once to logarithmic calculation, we can put the above still more plainly, thus:—

When extremes are conjunct.

86. RULE *a.* To find log sine middle { From log tang extr conjunct + log tang
other extr conjunct subtr rad (or 10)

87. RULE *b.* To find log tang extr conjunct { From log rad + log sine of middle
subtr log tang other extr conjunct

When extremes are disjunct.

88. RULE *c.* To find log sine middle { From log cos extr disjunct + log cos
other extr disjunct subtr rad


89. RULE *d.* To find log cos extr disjunct { From rad + log sine middle subtr
log cos other extr disjunct

care being taken to use complements of angles and hypotenuse. We will take an example, and at once apply the above formulæ.

90. In the right-angled triangle suppose the following:—

Given, a side $AC = 30^\circ$ } required, hypotenuse AB and $\angle A$.
 a side $BC = 40$ }

Fig. 52.



In constructing a spheric figure, it is, moreover, always convenient to place a given side on the primitive: we do so by making $AC = 30^\circ$ (by scale of chords) join CO .

By Problem V. draw a small circle parallel to the primitive $= 40^\circ$ distance; and where this cuts OC will be the third angular point B . Through the points ABD draw a great circle (by Prob. IV.), and ABC will be the triangle, of which the $\angle C$ will be a right angle (because the right circle CO passes through O , the pole of the primitive, and therefore is perpendicular to the arc AC), and its opposite side will be AB the hypotenuse.

N.B. It is convenient to mark the given parts as in the figure, and the parts required with an o .

To find the hypotenuse AB .

It will be seen that in order to get two "parts" alike (that is, two parts which shall be either *conjunct* or *disjunct*), the middle part should in the above figure be the hypotenuse itself, for it is separated from the parts given by

$\angle A$ at one end, and by $\angle B$ at the other. The parts AC and BC are therefore called extremes *disjunct*.

91. The following rhymes may help the memory in applying tangents or cosines, &c. :—

Tangents join the middle
(Put the middle where you please);
Cosines afar,
From middle are
(Five parts you have in these).

We have, therefore, in this example, to use *cosines* with the extremes (subject to the correction for hypotenuse and angles), and we want to find the middle part, AB . Rule C gives the following equation :—

$$\text{Sine of middle part} = \frac{\cos \text{ ext. disj.} \times \cos \text{ of other ext. disj.}}{\text{radius.}}$$

Now, before proceeding, let us consider what is meant by this equation, and why it was further altered into rule c (88). We have already shown that multiplication is performed in logarithms by addition; and division by subtraction; and in the fraction standing on the right side of the sign of equation, there are two quantities to be multiplied, and a quantity which is to divide their product. We therefore *add* the logarithms of the two factors in the numerator from the sum and *subtract* the logarithm of the denominator.

92. In a proportion worked by logarithms it is better to place the terms vertically (putting the divisor as the first term), thus :—

As radius
is to cos extreme disjunct,
so is cos of the other extreme disjunct
to the sine of the middle part.

(Remember old Dr. Kelly's Hibernian rhyme :—)

“Now the product of radius and middle part sine,
Equals that of the tangents of parts that combine,
And also the cosines of those that disjoin.”)

93. But we have first (83) to correct the above proportion if the hypotenuse or an angle form part of it. It should therefore appear accurately thus:—

as radius	.	.	.	co ar	10·000000
is to cos side A C 30°	9·937531
so is cos side B C 40°	9·884254
to cos hyp. A B	<u>9·821785</u> = 48° 26' 21"

To find $\angle A$:—Use the rule of sines (79), (having now opposite sides and angles).

as sine A B 48° 26' 21"	.	.	.	co ar	0·125952
to sine of opp. \angle (radius)	10·
so is sine of B C 40°	9·808067
to sine of $\angle A$	<u>9·934019</u> = 59° 12' 37"

But as it is better to thoroughly master one question in all its bearings, we will take a different view of the same question, and determine on finding the angle A, before we find the hypotenuse.

We now evidently call A C the middle, and then the $\angle A$ and the side B C will be extremes *conjunct* (the right angle is not one of, and *does not separate* the parts remember), and Rule 87 gives us as an equation:—

To find log tang extr conjunct $\left\{ \begin{array}{l} \text{From log rad + log sine of middle subtr} \\ \text{log tang of other extr conjunct} \end{array} \right.$

tang of the other ext. conj. B C 40°	.	.	.	co. ar.	0·076186
rad	10·
sine of middle A C 30°	9·698970
co tang $\angle A$	<u>9·775156</u> = 59° 12' 37"

The hypotenuse can now be found by the rule of sines (78), thus:—

as sine $\angle A$ 59° 12' 37"	.	.	.	co ar	0·065981
to sine side B C 40°	9·808067
sine 90°	10·
sine hyp.	<u>9·874048</u> = 48° 26' 21"

94. The co. ar. (read arithmetical complement) of an arc is what the logarithm wants of radius, and is readily formed by subtracting each figure of the logarithm (*beginning at the left hand*) from 9 and the last from 10: thus the logarithmic co. ar. of $\cdot 333333$ is $\cdot 666667$. This saves subtraction as the three logs may then be added.

Every right-angled spheric triangle may in like manner be worked by the four equations A, B, C, D, or a, b, c, d (page 65). But in such triangles as have a side for a right angle, a modification of the above rules is necessary, for we have in such cases what are called

QUADRANTAL SPHERIC TRIANGLES.

95. The only difference in the mode of working arises from an apparently whimsical perversion of terms.

For now the merry *quadrant*
 Its pranks with us to play,
Transforms itself to radius,
 And laughs our rules away.
 It calls legs, angles!—angles, legs!
 Our notions to confuse;
 While its opposite angle's supplement,
 It calls hypotenuse!

So that, considering the quadrantal side as radius, and the supplement of its opposite angle as hypotenuse, the solution of quadrantal angles is performed by rules already explained; viz., the rule of sines, and the four rules for the five circular parts.

OBLIQUE SPHERIC TRIGONOMETRY.

96. The majority of spheric questions which occur in practice fall under the denomination “oblique,” *i. e.* having neither an angle nor a side equal to a right angle.

97. Oblique spheric trigonometry admits of the six following cases, viz.:—

The given parts will be either

1. Two sides and an opposite angle.
2. Two angles and an opposite side.
3. Two sides and an included angle.
4. Two angles and an included side.
5. Three sides.
6. Three angles.

98. The solution of the first four cases may either be effected by means of a perpendicular let fall from one of the angles to its opposite side, or by special rules not requiring the perpendicular. There is a little difficulty with beginners in constructing the triangle so as to admit of a perpendicular being drawn in *such manner as to retain two of the given parts in one of two new right-angled triangles thus formed*. This may often be avoided by attending to the following directions, viz., describe the usual circle and quarter it. Then lay off a given side, A C, on the primitive (A being the angular point of the *left* hand of the side of the triangle, which lies on the primitive, and C the other end of it. It is merely convenient to have one method), and at C lay off the given angle (Prob. XI). Consider how you can secure two of the given parts in a new right angle you are about to form, and (by Prob. X.) let fall the perpendicular accordingly.

99. When the two angles which lie upon the side which is to be crossed by a perpendicular are of *like affection*, i. e., both greater or both less than a right angle, the perpendicular will fall *within* the triangle, but when unlike, *i. e.* when one is *acute* and one *obtuse*, the perpendicular will fall *without* the triangle.

100. N.B. In explaining the nature of "construction" from observation, it was recommended that the various

circles of the sphere should always be made to represent their respective parts in the general astronomic diagram, but for calculation it is better to take the parts merely as sides or angles.

The following will show the method of "projecting" and calculating any oblique spheric triangle which can possibly occur.

101. CASE I.


Two sides and an opposite angle.

Given, a side 60°
 „ a side 100°
 „ an opposite angle 130° } To find a side and the other angles.

By construction :—

N.B. In nearly all cases we suppose a circle to be already drawn with a chord of 60° , and two diameters at right angles within it.

Fig. 53.



Lay off from C to A the one side, 60° (from scale of chords). From A draw A G, with the supplement of 130° , and draw (by Prob. VIII.) the great circle A B G. About E

describe the small circle vy , at a distance of the supplement of 100° (or 80°) from it (by Prob. VI.); and where vy cuts ABG is the angular point B .

With the three points $EB C$ draw an oblique circle (by Prob. IV.), and ABC is the triangle.

To measure the parts required, viz., AB , $\angle C$, and $\angle B$:

AB is measured by Prob. XII.

$\angle C$ is measured by Prob. XIII.

$\angle B$ is measured by Prob. XIV.

By calculation :—

1st. *By means of a Perpendicular.*

Having a side on the primitive with an adjacent $\angle A$ given, let fall a perpendicular from the $\angle C$ upon AB (produced if necessary) to D (Prob. X.).

To find the other opposite $\angle ABC$ by rule of sines (79).

as sine $BC\ 100^\circ$	co ar	0.006649	
to sine $\angle A\ 130^\circ$:	9.884254	
so is sine $AC\ 60^\circ$:	9.937531	
to sine $\angle ABC$:	9.828434	= $42^\circ\ 21'$
		<u>180</u>	
			$\angle CBD = 137\ 39$

To find AD (in triangle ADC):—

The $\angle A$ will be the middle part, and the hypotenuse AC and the side AD will be extremes conjunct (87).

as cotang hyp $AC\ 60^\circ$	co ar	0.238561	
to rad	:	10.	
cos middle $\angle A\ 130^\circ$:	9.808067	
to tang	:	10.046628	= $48^\circ\ 4'$
		<u>180</u>	
			$\text{tang } AD = 131\ 56$

To find DB (in triangle CBD):—

The $\angle B$ will be middle, and hyp BC and the side DB will be extremes conjunct (87).

as tang BC 100°	co ar	0.753681	
to radius		$10.$	
cos $\angle B$ $137^\circ 39'$		9.868670	
tang DB		10.622351	$= 76^\circ 35'$

To find $\angle C$ by rule of sines (79).

$AD = 131^\circ 56'$	as sine AC 60°	. . .	0.062469
$DB = \frac{76}{37}$	to sine $\angle B$ $42^\circ 21'$		9.828434
$BA = \frac{55}{19}$	so is sine BA 55 19		9.915035
	to sine $\angle C$		$9.805938 = 39^\circ 46'$

2nd. Without a Perpendicular.

When the solving of a spheric triangle presents any difficulties to the unpractised as to where to let fall the perpendicular, if time is precious, recourse may be had to the following rule:—

When two sides and an opposite angle are given.

First find the angle opposite to the other of the two given sides, and the third angle of the triangle may then be found thus:—

RULE.—As the sine of half the difference of the two sides is to the sine of half their sum so is the tangent of half the difference of the two angles to the cotangent of half the contained angle.

Or, as applied to the preceding question, viz. :—

Given, a side AC =	60°
a side BC =	100°
$\angle A$	$= 130^\circ$
$\angle B$	$= 42^\circ 21'$

To find the angle C:—

side AC	60°	as sine half diff 20° . . .	co ar	0.465948
side BC	100	to sine half sum 80° . . .		9.993351
	<u>160</u>	tang half diff angles 43° 49½' . . .		9.982182
half sum	<u>80</u>	cotang half contd ∠ C 19° 53' =		<u>10.441481</u>
half diff	20	∴ whole ∠ C is 39° 46' . . .		
∠ A 130°				
∠ B 42 21				
	<u>87 39</u>			
half diff	43 49½			

Then the side AB may be found by the rule of sines.

102. CASE II.


Two angles and an opposite side.

Given, an angle 60°	}	To find a side
angle 70°		a side
opp. side 50°		an angle

By construction:—

At the point C make an angle of 60° with the primitive (Prob. VIII.) by drawing CBE; about C, at the distance of

Fig. 54.



50° (the given side), draw the small circle Ay (by Prob. IX.); at B, where these intersect, draw (by Prob. XI.) the great circle AB, making an angle of 70° with the primitive, and ABC will be the spheric triangle.

To measure the parts required, viz. AC, AB, and $\angle B$:—

1. measure AC on the scale of chords = $49^{\circ} 37'$
2. AB is measured by Prob. XII. . = $44 55$
3. $\angle B$ is measured by Prob. XIV. . = $110 52$

By calculation :—

1st. *By means of a Perpendicular.*

In order to preserve the two parts BC and $\angle C$ in the same rectangle, let fall a perpendicular on side AC. This is done by drawing a line from the centre of the circle, which is always the pole of the primitive, through the $\angle B$ till it cuts AC at D. (Prob. X.)

To find the other opposite side AB by rule of sines :—

as sine $\angle A 70^{\circ}$. co ar	0.027014	
to sine side BC 50°	. .	9.884254	
so is sine $\angle C 60$. .	<u>9.937531</u>	
to sine side AB	. .	9.848799	= $44^{\circ} 54' 35''$

To find the segment CD in $\triangle BCD$:—

The angle C will be “middle,” and the hypotenuse BC and CD will be extremes conjunct (87).

as cotang hyp BC 50°	. co ar	0.076186	
to rad	10.	
so is cosine $\angle C 60^{\circ}$. .	<u>9.698970</u>	
to tang side CD	. .	9.775156	= $30^{\circ} 47' 23''$

To find segment AD in triangle ABD :—

The angle will be *middle*, and the side AD and hyp. AB will be extremes conjunct (87).

as cotang AB $44^{\circ} 54' 35''$	co ar	9.998631	
to radius	10.	
so is cos $\angle A 70^{\circ}$. .	<u>9.534052</u>	
to tang segment AD	. .	9.532683	= $18^{\circ} 49' 36''$
seg CD = $30^{\circ} 47' 23''$			
seg AD = <u>18 49 36</u>			
49 36 59			= side AC

To find $\angle B$ in $\triangle ABC$ by rule of sines (79).

as sine side BC	= 50°	. co ar	0.115746
to sine $\angle A$	= 70	.	9.972986
so is sine side AC	= 4937	.	<u>9.881799</u>
to sine $\angle B$			9.970531 = 69° 7' 51"
			180
			$\therefore \angle B = 110\ 52\ 9$

N.B. We take the supplement of 69° 7' 51" because the construction shows the $\angle B$ to be obtuse.

2nd. Without a Perpendicular.

The side AB opposite the other given angle can be found by rule of sines as before, and equals 44° 54' 35". Then find $\angle B$ by the special rule given in Case I. thus:—

a side AB	44° 54' 35"		$\angle A$	70°
a side BC	50		$\angle C$	60
	2)94 54 35			2)10
half sum of sides	47 27 18	half diff two angles		5
diff two sides	5 5 25			
half diff two sides	2 32 42			

as sine $\frac{1}{2}$ diff 2 sides	2° 32' 42"	co ar	1.352577
to sine $\frac{1}{2}$ their sum	47 27 18	.	9.867318
so is tang $\frac{1}{2}$ diff 2 angles	5 0 0	.	<u>8.941952</u>
to cotang $\frac{1}{2}$ contained $\angle B$			10.161847 = 34 33 46
			2
			69 7 32
			180
			$\angle B = 110\ 52\ 28$

Find AC by rule of sines (78).

as sine $\angle A$ 70°	. . .	co ar	0.027014
to sine side BC 50°	9.884254
so is sine $\angle B$ 110° 52' 28"	<u>9.970516</u>
to sine side AC	. . .	=	9.881784 = 49° 36' 51"

103. CASE III.

Two sides and an included angle.

Given, a side AC = 60°	}	To find side BC.
„ a side AB = 110°		„ $\angle B$
„ the included angle A 45°		„ $\angle C$

By construction :—


Lay off $AC = 60^\circ$ from the scale of chords.

Draw the great circle ADB (by Prob. XI.), at the distance of 45° from the primitive; about G draw the parallel circle vy , distance of the supplement of 110 (Prob. V.); where the two circles cross is the point B , through the points $C B H$ draw a great circle (Prob. IV:), and ABC will be the triangle.

To measure the parts required:—

1. The $\angle C$, ax is measured on the scale of semi-tangents (Prob. XIII.).
2. measure $\angle B$ by Prob. XIV.
3. measure side CB by Prob. XII.

Fig. 55.



By calculation :—

1. *By means of a Perpendicular.*

Draw a perpendicular from the $\angle C$ upon the side AB to D (by Prob. X.).

Find the segment AD in the $\triangle ACD$ (78).

Then the $\angle A$ is the middle, and hyp AC and side AD are extremes conjunct (87).

as cotang hyp $AC\ 60^\circ$	co ar	10·238561
to radius		10·
so is cos $\angle A\ 45^\circ$		9·849485
to tang of segment AD	$50^\circ 46' 17'' =$	10·088046
subtr from side AB	110	
segment		<u>59 13 43</u>	= segment AD

Find DC in $\triangle ADC$ by rule of sines (78):—

as rad.	co ar	10·000000
sine side (hyp.) $AC\ 60^\circ$		9·937531
sine $\angle A\ 45^\circ$		9·849485
to sine side DC	<u>9·787016</u>	= $37^\circ 46'$

Find $\angle B$ in $\triangle BDC$.

Here BD is middle, and DC and $\angle B$ are extremes conjunct.

as tang $DC\ 37^\circ 46'$	co ar	0·110839
to rad.		10·
so is sine side $BD\ 59^\circ 14'$		9·934123
to cotang $\angle B\ 42^\circ 2'$	<u>10·044962</u>	

Find side BC by rule of sines (78):—

as sine $\angle B\ 42^\circ 2'$	co ar	0·174209
to sine $AC\ 60^\circ$		9·937531
so is sine $\angle A\ 45^\circ$		9·849485
to sine side BC	<u>9·961225</u>	= $66^\circ 9'$

Find $\angle C$ by rule of sines (79):—

as sine side $AC\ 60^\circ$		0·062469
to sine $\angle B\ 42^\circ 2'$		9·825791
so is sine side $AB\ 110^\circ$		9·972986
to sine $\angle C\ 46^\circ 36'$	<u>9·861246</u>	

180

$$\angle C = 133^\circ 24'$$

N.B. We take the supplement because the $\angle C$ is obtuse (by construction).

2. *Without a Perpendicular.*

RULE.—When two sides and an included angle are given:—

1. As the sine of half the sum of the two sides is to the sine of half their difference so is the cotangent of half their contained angle to the tangent of half the difference of the other angles;

and again,

2. As the cosine of half the sum of the two sides is to the cosine of half their difference, so is the cotangent of half the contained angle to the tangent of half the sum of the other two angles.

And half the difference thus found added to half the sum gives the greater angle; and half the difference subtracted from the half sum gives the smaller angle.

side A C = 60°

side A B = 110

$$\begin{array}{r} 2)170 \\ \hline \end{array}$$

half sum $\frac{85}{}$

$\angle A$ (contained angle) = 45°

$$\begin{array}{r} 2)50 \\ \hline \end{array}$$

half diff $\frac{25}{}$

Then by the above rules:—

as sine $\frac{1}{2}$ sine of 2 sides 85°	co ar	0.001656	
to sine $\frac{1}{2}$ diff ditto 25°		9.625948	
so is cotang $\frac{1}{2}$ contained $\angle A$ 22° 30'		10.382776	
to tang $\frac{1}{2}$ diff of other angles		10.010380	= 45° 41'
as cos $\frac{1}{2}$ sum 85°	co ar	1.059704	
to cos. $\frac{1}{2}$ diff. 25°		9.957276	
so is cotang of $\frac{1}{2}$ the cont. $\angle A$ 22° 30'		10.382776	
to tang $\frac{1}{2}$ sum of other angles		11.399756	= 87° 43'
				greater $\angle C$ = 133 24
				less $\angle B$ = 42 2

Find BC by rule of sines (78).


as sine $\angle C$ $133^{\circ} 24'$	co ar 0.138720
to sine side AB 110°	9.972986
so is sine $\angle A$ 45°	9.849485
	9.961191 = $66^{\circ} 8'$

. 104. CASE IV.

Two angles and an included side.

Given, an angle 60°	}	To find an angle
,, an angle 36°		a side
,, included side 70°		a side.

Fig. 56.



By construction :—

Lay off AC equal to 70° from the scale of chords.

Draw the great circle AB (Prob. XI.), at a distance of 36° from the primitive, and in like manner CB at a distance of 60° ; where these two intersect will be the point B, and ABC will be the triangle.

To measure the parts required :—

1. measure side AB by Prob. XII.
2. measure side BC ,, XII.
3. measure $\angle B$,, XIV.

By calculation :—

1. *By means of a Perpendicular.*

Let fall a perpendicular from $\angle A$ on side BC produced (Prob. X.) (or from $\angle C$ on AB produced, suppose the former).

Then will ADC be a right angle, as will also ADB right angled at D .

Find the $\angle DAC$ in the $\triangle DAC$.

The side AC will be middle, and the angle C , and $\angle DAC$ will be extremes conjunct (82): then by (87):—

as cotang $\angle C$ 60°	co ar	0.238561
to rad	10.
so is cosine side AC 70°	9.534052
to cotang $\angle DAC$	9.772613 = $59^\circ 21'$
						<u>$\angle BAC - 36$</u>
						$\angle DAB = 23 \ 21$

Find AD in $\triangle ACD$.

The angle DAC is middle, and extremes are conjunct.

as cotang side $AC = 70^\circ$	co ar	0.438934
to rad	10.
so is cos $\angle DAC$ $59^\circ 21'$	9.707393
to tang AD	10.146327 = $54^\circ 28'$

Find AB in $\triangle ADB$.

The $\angle DAB$ is middle, and sides AD and AB are extremes conjunct.

*as tang side AD $54^\circ 28'$	co ar	9.853673
to rad	10.
so is cos $\angle DAB$ $23^\circ 21'$	9.962890
to tang side AB	9.816563 = $56^\circ 45'$

* It promotes accuracy to use the logarithm itself, from which the tang AD was taken on the previous calculation. 9.853673 is the ar. co. of 10.146327.

Find BC by rule of sines (78).

as sine $\angle C$ 60°	. . .	co ar	0.062469
to sine side AB $56^\circ 45'$. . .		9.922355
so is sine $\angle A$ 36°	. . .		<u>9.769219</u>
to sine side BC . . .			9.754043 = $34^\circ 35'$

Find $\angle ABC$ by rule of sines (79).

as sine side AB $56^\circ 45'$	co ar	0.077645
to sine $\angle C$ 60°	. . .	9.937531
so is sine side AC 70°	. . .	<u>9.972986</u>
to sine $\angle ABC$. . .		= 9.988162 = $76^\circ 41'$
		<u>180 0</u>
	$\angle B$	103 19

N.B.—The supplement is used because construction shows the angle to be obtuse.

2. Without a Perpendicular.

RULE.—When two angles and the included side are given:—

As the sine of half the sum of the two angles
is to the sum of half the difference
so is the tangent of half the contained side
to the tangent of half the difference of the other two sides.

And again:—

As the cosine of half the sum of the two angles
is to the cosine of half their difference,
so is the tangent of half the included side
to the tangent of half the sum of the other sides;

and half the difference added to half the sum will give the greater side, and half the difference subtracted from half the sum will give the smaller side.

an angle	36°	side	$\frac{70^\circ}{2} = 35^\circ =$ half the included side.
an angle	<u>60</u>		
	2)96		
half sum	<u>48</u>		
	2)24		
half diff	<u>12</u>		

Find sides AB and BC.

as sine $\frac{1}{2}$ sum 2 angles 48° co ar 0.128927
to sine $\frac{1}{2}$ diff 12° 9.317879
so is tang $\frac{1}{2}$ included side 35° 9.845227
to tang $\frac{1}{2}$ diff other sides <u>9.292033 = $11^\circ 5'$</u>
as cos $\frac{1}{2}$ sum of 2 angles 48° co ar 0.174489
to cos $\frac{1}{2}$ diff 12° 9.990404
sine tang $\frac{1}{2}$ included side 35° 9.845227
to tang $\frac{1}{2}$ sum other sides <u>10.010120 = $45^\circ 40'$</u>
	greater side AB = <u>$56 45$</u>
	smaller side BC = <u>$34 35$</u>

Find $\angle B$ by rule of sines (79).

as sine side EC $34^\circ 35'$ co ar 0.245954
to sine $\angle 36^\circ$ 9.769219
so is sine side AC 70° <u>9.972986</u>
to sine $\angle B$ <u>9.988159 = $76^\circ 41'$</u>
	<u>180</u>
	$\angle B$ <u>$130 19$</u>


N.B.—The construction of the figure shows the $\angle B$ to be obtuse; therefore we use the supplement.

105. CASE V.

Three sides given:—

- side AC 60°
- side AB 70°
- side BC 100°

Fig. 57.



By construction:—

Lay off one side (say 60°) on the primitive, as at AC, from chord of 60° . About E draw (Prob. VI.) a small circle *ay* at the distance of the supplement of BC. About A draw a small circle *xz* at the distance of 70° (Prob. VI.). Through point B, where the small circles intersect, draw (by Prob. IV.) an oblique circle AB, and ABC will be the triangle.

To measure the parts required:—

1. *mn* on the scale of semi-tangents measures $\angle A$.
2. The $\angle C$ is measured from Z to *o* (semi-tangent backwards from 90°).
3. The $\angle B$ by Prob. XIV.

By calculation:—

RULE.—Find half the sum of the three sides. Subtract from this half sum each of the two sides which, together, contain the required angle. Then add the sines of these two remainders to the sines of the two sides which contain the angle (using the co arcs of the latter). Half the sum of these four logarithms will give the sine of half the required angle.

To find the angle A:—

side AC = 60°	sine	co ar	0.062469
side AB = 70°	sine	co ar	0.027014
side BC = 100°			
			<u>2)230</u>
half sum of sides			115
side AC = 60			<u>55</u>
		first remainder	9.913365
			<u>115</u>
side AB = 70			<u>45</u>
		second remainder	9.849485
			<u>2)19.852333</u>
	sine	$57^\circ 32'$	= 9.926166
			<u>2</u>
$\angle A =$			<u>115 4</u>

or 50° , as ef , from scale of semi-tangents. Where the two small circles cut, as at n , will be the pole of the oblique circle, which shall make an angle of 45° with the primitive and 50° with the other oblique circle.

Through on draw a diameter; measure on on the scale of semi-tangents, and lay off its complement beyond o ; say to v .

Through the three points DvC (DC being at right angles to GH) draw the oblique circle DvC (Prob. IV.), and ABC shall be the triangle.

To measure the parts required:—

1. AC is measured on the scale of chords.
2. AB „ by Prob. XII.
3. BC „ by Prob. XII.

By calculation:—

RULE.—From half the sum of the three angles take each of the angles next to the side required. Add the co arcs of the sines of the two angles which adjoin the required sides, together with the cosines of the two remainders. Then half the sum of these logarithms will equal half the cosine of the side required.

Find side BC .

$\angle A =$	130°			
$\angle B =$	50	sine . . .	co ar	0.115746
$\angle C =$	45	sine . . .	co ar	0.150515
	$2)225$			
	$112\frac{1}{2}$			
$\angle B =$	50			
	$62\frac{1}{2}$	1st remainder co sine		9.664406
	$112\frac{1}{2}$			
$\angle C =$	45			
	$67\frac{1}{2}$	2nd remainder co sine		9.582840
				$2)19.513507$
				9.756753
		cosine $34^\circ 50'$	=	
		2		
		$69 40$		
		180		
		$110 20$	= side BC	

Find side AC by rule of sines (78).

as sine $\angle A$ 130°	. . .	co ar 0.115746
to sine BC $110^\circ 20'$. . .	9.972986
so is sine $\angle B$ 50°	. . .	<u>9.884254</u>
to sine side AC	. . .	$9.972986 = \angle AC 70^\circ$

Find side AB (78).

as sine $\angle A$ 130°	. . .	co ar 0.115746
to sine BC $110^\circ 20'$. . .	9.972986
so is sine $\angle C$ 45°	. . .	<u>9.849485</u>
to side AB	. . .	$9.938217 = \angle AB 60^\circ 9'$

107. All the rules necessary for calculating a spheric triangle have been explained. Other formulæ might have been added, but where the application of the subject to practice is the main object sufficient has been given. But in order to obviate any possible difficulties which a student might at first encounter in the application of what has been said, we shall now return to the examples which were constructed from supposed observation (page 65), and show the manner of calculating the results; and the more especially is this necessary, because, in spherics, the triangle drawn as a problem in nautical astronomy differs from that which is more adapted to calculation; and again this, if an oblique triangle, requires some management in order to so construct the triangle as to render it convenient for letting fall an available perpendicular to be used with the "5 circular parts," or Napier's rules. An example (No. 2) will be given in illustration of this difference, while the others will be worked without a perpendicular, leaving it to the student to exercise his ingenuity in further construction or calculation.

108. EXAMPLE 1.—To find the latitude of a place:—

Given, meridian altitude 60° (The observer north of the sun)
 ,, declination . 20° N.

This question needs no other figure than that already

Find $\angle C$ by rule of sines (79).

as sine side BC 70°	co ar	9.027014
to sine $\angle A$ 135°		9.849485
so is sine side AB 40°		9.808067
to sine $\angle C$		<u>9.684566</u> = $28^\circ 56'$

side 70	\angle $135^\circ\text{-}0'$
side 40	\angle $28^\circ\text{-}56'$

2)110	2)106.4
-------	---------

$\frac{1}{2}$ sum . . . 55 $53.2 = \frac{1}{2}$ diff \angle s.

2)30

$\frac{1}{2}$ diff . . . 15

as sine $\frac{1}{2}$ diff of sides 15°	co ar	0.587004
to sine $\frac{1}{2}$ sum 55°		9.913365
so is tang $\frac{1}{2}$ diff $53^\circ 2'$		<u>10.123411</u>
to tang $\frac{1}{2}$ contained \angle		<u>10.623780</u> = $13^\circ\text{-}23'$


2
B = $26^\circ 46'$

Find side AC by rule of sines.

as sine $\angle C$ $28^\circ 56'$	co ar	0.315342
to sine side AB 40°		9.808067
so is sine $\angle B$ $26^\circ 46'$		<u>9.653558</u>
to side AC		<u>9.776967</u> = $36^\circ 45'$

90
latitude = $53^\circ 15'$

Fig. 60.



N.B.—Notice that while CR will be the latitude, we, in forming the triangle ABC, use *the complement* of the two given sides, and that the side required in the triangle ABC is therefore the co latitude.

2. By means of a Perpendicular.


As previously recommended (90), it is well, as a general rule, to put a given angle at C (by Prob. XI.), and a given side upon A C by scale of chords, letting fall the perpendicular upon B C (or B C produced if necessary) from the angle A by Prob. X.

In this question we have two sides, and their two opposite angles, one of the latter having been found by rule of sines.

The following will be the triangle as constructed for the perpendicular let fall from A :—

side A C being	70
side A B „	40
$\angle C$ „	28 56
$\angle B$ „	135

Fig. 61.



Find D C in $\triangle A D C$.

The $\angle C$ is middle and extremes are conjunct (82).

Then correcting for complements or supplements (83), we have by (87)

as cotang A C 70°	co ar	0.438934
rad		10.
cosine $\angle C$ $28^\circ 56'$		9.942099
to tang side D C		<u>10.381033</u> = $67^\circ 25'$

Find AD in $\triangle ADC$.

AD is middle, and extremes are disjunct, then by (88).

as rad	10
to sine AC 70°	9.972986
so is sine $\angle C$ $28^\circ 56'$	9.684658
to sine AD	<u>9.657644</u> = $27^\circ 2'$

Find DB in $\triangle ADB$.

AB is middle and disjunct, then by (89).

as cos AD $27^\circ 2'$	0.050248
to rad	10
so is cos AB 40°	9.884254
to side DB	<u>9.934502</u> = $30^\circ 41'$
									side DC <u>67 25</u>
									co lat or side BC = <u>36 44</u>
									90
									<u>latitude required = $53^\circ 16'$</u>

It would have been shorter (perhaps not so obvious to a beginner) to have found DB by using the $\triangle ADB$, and using the complement of ABC (= 45°) as the $\angle DBA$, thus:—

The $\angle B$ 45° is middle and conjunct, then by (87).

as cotang 40°	9.923814
to rad	10
so is cos $\angle B$ 45°	9.849485
to tang DB	<u>9.773299</u> = $30^\circ 41'$ co lat

110. EXAMPLE 3.—To find the latitude of a place :

Given altitude 50° south of the sun
 dec 10 S.
 time 10 A.M.

In this example, the sun being in south declination and the observer being south of the sun, it is evident that the observer must be in south latitude (remember that latitude is the height of the pole of the sphere above the horizon).


By construction :—

Being in south latitude the south pole will be above the south part of the horizon. We generally, in construction, consider the north part of the horizon to lie on the right of the centre of the figure, and the south part on the left part.

Assume S the south pole, and draw SOP , and EQ , the equator at right angles to it.

Draw the great circle SxP (by Prob. XI.), making an angle of two hours or 30° with primitive.

By Prob. VI., draw the parallel of declination dc at a distance of 10° south of EQ , or 80° from the primitive.



To draw the great circle ZxN , so that $Zx = 40^\circ$ (the co of 50° the altitude), take the secant of 40 from the centre o , and sweep an arc as gn , and the tangent of 40° from the point x , and sweep another arc hm across it; the intersection will be the zenith line OZa (Prob. VI.). With the three points ZxN , describe the great circle ZxN (Prob. IV.), and SZx is the triangle, and SZ the co latitude, SH the latitude required.

By calculation :—

Find $\angle Z$, by rule of sines (79).

as sine of side ZX = 40° . . .	co ar	0.191933	
is to sineside of $\angle S = 30$. . .		9.698970	
so is sine side Sx = 80 . . .		9.993351	
		9.884254	= 50°
to sine $\angle Z$ = . . .			180°
(obtuse by construction)			130° = $\angle Z$

To find $\angle x$.

a side Zx = 40° . . .	40°	an angle	130°	
a side Sx = 80 . . .	80	an angle	30	
	2)120		2)100	
half sum sides = 60	$\frac{1}{2}$ diff 20	50	half diff 2 angles	

By rule (page 73):—

as sine $\frac{1}{2}$ diff 2 sides 20° . . .	co ar	0.465948	
is to sine $\frac{1}{2}$ their sum 60° . . .		9.937531	
so is tang $\frac{1}{2}$ diff 2 angles 50° . . .		10.076186	
to co tang $\frac{1}{2}$ then contained \angle . . .		10.479665	= 18° 20'
			$\angle x = 36° 40'$

To find co lat Sz, by rule of sines (78).

as sine $\angle S = 30°$. . .	co ar	0.301030	
is to sine side Zx = 40° . . .		9.808067	
so is sine $\angle x = 36° 40'$. . .		9.776090	
to sine side SZ . . .		9.885187	= co lat 50° 9' S.
			90
			lat 39° 51'


111. EXAMPLE 4.— To find the latitude by a celestial body on the meridian below the pole :

Given, altitude of α Lyræ on meridian below the north pole = 20°
 declination of $\delta\gamma$ = 38½ N.

This forms no triangle, the star being on the primitive.

Let HR be the horizon. Lay off the alt 20° to a , the star's place. The star being $38\frac{1}{2}^\circ$ N. of the equator (in declination), lay off $38\frac{1}{2}^\circ$ from a to Q. Draw QE and PS, and PR will be the latitude, and will equal $90^\circ - (38\frac{1}{2}^\circ - 20^\circ) = 90^\circ - 18\frac{1}{2}^\circ = 71\frac{1}{2}^\circ$ north.


Fig. 63.



112. EXAMPLE 5.—To find the latitude of a place :—

Given, app. time 9 a.m.
 declination, 20° N.
 azimuth, S. 60° E.
 Observer N. of sun.

Fig. 64.



By construction :—

Assume a point P on the primitive, and draw PS and EQ. Draw the oblique circle Px, making 45° with the primitive (Prob. XI.). About P draw the parallel circle dx 70° (the co declination) distant from P (Prob. V.). Through the intersection x draw Zx, making the angle at the primitive = 60, the azimuth from noon, or south (Prob. XI.). Then ZxP will be the spheric triangle, and ZP the co latitude.

By calculation :—

To find Zx by rule of sines (78).

	60° as sine ∠Z 120° . . .	co ar 0·062469
	186 is to sine side xP 70° . . .	9·972986
∠x = 120	so is sine ∠P 45° . . .	9·849485
	to sine side Zx . . .	9·884940 = 50° 6'

To find ∠x (page 73).

sine	50° 6'	∠	120°
side	70	∠	45
	<u>2)120 6</u>		<u>2)75</u>
half sum	60 3		37 30 half diff
	<u>2)19 54</u>		
half diff	9 57		
as sine ½ diff 2 sides	9° 57' . . .	co ar	0·762485
to sine ½ sum 2 sides	30° 3' . . .		9·937749
so is tang ½ diff 2 angles	37° 30' . . .		9·884980
to cotang ½ cont ∠x . . .			10·585214 = 14 34'
			<u>2</u>
		∠x	29 8

Find side ZP by rule of sines (78).

as sine ∠Z 120° . . .	co ar 0·062469
is to sine side Px 70° . . .	9·972986
so is sine ∠x 29° 8' . . .	9·687389
to sine of side ZP . . .	9·722844 = 31° 53'
	<u>90</u>
	latitude ZP 58 7 N

side . . .	80° 0	∠ 135° 0 0'	
side . . .	44 20	∠ 30 7 0	
	2)124 20	2)104 53 -0	
half sum . . .	62 10	52 26 30	half diff ∠ s
	2)35 40		
half diff . . .	17 50		

Find ∠ P (page 73).

as sine of half diff sides 17° 50'	. . . co ar 0·513925'
to sine of half sum 62° 10' 9·946604
so is tang half diff ∠ s 52° 26½' 10·114104
to cotang half cont ∠ P 10·574633 = 14° 55'
	2
	hour angle ∠ P = 29 50

N.B.—To convert space into time and the reverse, use the following rules:—

RULE.—Multiply space by 4 and divide the degrees by 60, thus:—

the arc	29° 50'	
	4	
	60)119 20	
		1h 50m 20s = the above hour angle

To convert time into space:—

RULE.—Reduce hours to minutes and divide by 4, thus:—

the time	1h 59m 20s	
	60	
	4)119 20	
		29° 50'

114. EXAMPLE 7.—To find time.


Given, declination, 20° S. (being in N. lat.)
 altitude 20
 azimuth S. 45° W.

By construction:—

From Z draw ZxN, making an angle 45° with the primitive (Prob. VIII.). About Z draw the parallel ef at a

(considering P, as usual, the north pole of the world). About S, the south pole, draw a parallel equal to sun's north polar distance, or 110° (Prob VI.). About Z, the zenith, draw a parallel fg equal to co alt (or zenith distance) (Prob. VI.). Through ZxN draw an oblique circle (Prob. IV.) and ZxP will be the triangle and $\angle P$ the hour angle.

Fig. 67.



By calculation :—

N.B.—Three sides are given ; find $\angle P$ (page 84).

a side Zx . = 60°
 a side ZP . = 69 co ar sine . 0.029848
 a side xP . = 110 co ar sine . 0.027014

2)239

half sum 119 30'
 ZP 69
50 30 = 1st remr . 9.887406

119 30

110

9 30 = 2nd remr . 9.217609

2)19.161877

sine $22^\circ 24' = 9.580939$

2

$\angle P$ 44 48

4

60)179 12

time from noon $2h\ 59m\ 12s$


117. EXAMPLE 10.—To find an azimuth.

Given, latitude	21° N.	declination	20 S	
time	9 a.m.		90	
declination	20° S.		<hr/>	110 = polar distance from N

By construction:—

Place the north pole at P, 21° above the horizon R. Draw diameters. Draw the oblique circle P α S with the \angle 45° from the primitive (45° = 3 hrs.) (Prob. VIII.). About S, the south pole, draw a parallel equal to the co declination (by Prob. VI.), and through the intersection x draw (Prob. VI.) Z x N and Z x P will be the triangle, and \angle Z the azimuth required.

Fig. 69.



By calculation:—

We have two sides given and an included angle.

a side xP	110		
a side ZP	69	included \angle 2)45°	
	<hr/>		
	2)179		22 30 half contained \angle
half sum	89 30		
	<hr/>		
	2)41		
half sum	20 30		

To find the other angles (page 79).


as sine $\frac{1}{2}$ sum 2 sides 89° 30'	co ar	0.000017
to sine $\frac{1}{2}$ diff 2 sides 20 30		9.544325
so is cot $\frac{1}{2}$ contd \angle 22 30		<hr/> 10.382776
to tang $\frac{1}{2}$ diff other angles		9.927119 = 40° 13'
as cos $\frac{1}{2}$ sum 2 sides 89° 30'	co ar	2.059158
to cot $\frac{1}{2}$ diff 2 sides 20° 30'		9.971588
so is cot $\frac{1}{2}$ contd \angle 22 30		<hr/> 10.382776
to tang $\frac{1}{2}$ sum other angles		12.413522 = 89° 47'
sum is the azimuth greater \angle Z		<hr/> 130 00
less \angle x		49 34

118. EXAMPLE 11. — To find a ship's course when sailing on a great circle, and the distance between port and port.

Given, ship's latitude in	50° N.
latitude of place bound to	10° N.
the difference of longitude between the two places	60° W.

The term great circle "course" is deceiving, inasmuch as no part of a circle is a straight line. A ship could not sail upon a great circle without constantly changing her course by compass; and, therefore, "great circle sailing" is positively impracticable, because a great circle "track" cuts no two meridians at an equal angle. The passage of a ship along a great circle track is evidently a series of courses *tangential* to it.

Fig. 70.



Suppose, in the above figure, AB to be a given great circle track; upon a ship's starting at *a* for the point B a compass course, if long continued, would take her considerably away from her proper track, and, we will say, place her at *a'*. It is evident, therefore, that the *shorter these compass courses are made, the more will the ship keep to her true course*, and the shorter will be the distance required to be sailed over. As an instance: A ship was 12 hours in going from *b* to *b'*, and 24 hours in going (at the same rate) from *c* to *c'*; we find that in these cases she would be leaving her great circle track altogether. It remains, therefore, to provide a *ready method* of finding how to steer by compass so as to depart *as little as possible* from

our proper track, which, of course, would be the nearest distance between the port left and port bound to.


The term *tangent sailing* is the only correct designation of this method, and was first suggested to the Astronomer Royal by the author in 1857.

From the above example to find the first or initial tangent course, we proceed thus—

By construction :—

Consider the diagram as a hemisphere drawn on the plane of a ship's meridian (*the ship being somewhere on the meridian*). PS would represent the poles, and EQ the equator. Let L represent the latitude, or the

Fig. 71.



ship's place, and l the latitude of port bound to, or $LQ = 50^\circ$ and $lQ = 10^\circ$. Draw diameters to point L , and draw PxS , making an angle of 60° (the difference of longitude) with the primitive (Prob. VIII.). Draw the parallel circle ll' , and through the point of intersection x draw the oblique circle LxN , Prob. X., and the $\angle xLQ$ will be the tangent course required; measured on rn (being 90° distant) it equals $63^\circ 17'$.

By calculation:—

We have here two sides and an included angle.

a side 80 included $\angle 2)60^\circ$
 a side 40 $\frac{30}{30} = \text{half contained } \angle$

2)120
 half sum 60

2)40
 half diff 20

To find the other angles (page 79.)

as sine $\frac{1}{2}$ sum of sides 60° co ar 0.062469
 to sine $\frac{1}{2}$ diff 20° 9.534052
 so is co tang $\frac{1}{2}$ contained angles 30° 10.238561
 to tang $\frac{1}{2}$ diff angles 9.835082 = $34^\circ 22'$

cos $\frac{1}{2}$ sum of sides 60° co ar 0.301030
 to cosine $\frac{1}{2}$ diff 20° 9.972986
 so is cotang $\frac{1}{2}$ contained angle 30° 10.238561
 to tang $\frac{1}{2}$ sum other angles 10.512577 = $72^\circ 55'$

$\angle L$ 107 17
180

the tangent course $\angle xLQ$ 72 43

or S 72 43 W

119. EXAMPLE 12.—To find the distance between port and port upon a great circle (in the above example).

Find side Lx by rule of sines (78).


as sine $\angle L$ (as above) $107^\circ 17'$ co ar 0.020066
 is to sine side 80° 9.993351
 so is sine $\angle P$ 60° 9.937531
 to sine side Lx 9.950948 = $63^\circ 17'$
60

distance to make good Lx 3797 miles.

Suppose that, three days afterwards, the ship was in lat, not 50° but 45° N, and diff of long between the two places was not 60° , but 50° , what would be her altered course?


Working as above would give the course about 67° , and the distance about 55° , or 3300 geographical miles.

N.B.—By Saxby's spherograph (a simple instrument which thoroughly illustrates nautical astronomy, and works any spheric triangle without calculation) a great circle, or tangent course, is easily obtained in five to ten seconds; being thus rendered more simple than even a Mercator's course.



120. EXAMPLE 13.—To find the “latitude of vertex” of a great circle track.

Given, latitude of the ship	36° S.
“ bound to	40° S.
difference of long.	110°



By construction :—

Make EL equal to lat 36° (from the scale of chords) south of the equator, and L will be the ship's place on her meridian. Draw diameters Ll and mn. About the south pole, S, draw a parallel circle Lo, distant 50° the co lat (Prob. V.). Draw the oblique circle SxN, making 110° with the primitive (Prob. VIII.). Through the intersection x draw Lx l, and LxS will be the triangle. Let fall a perpendicular Sy upon Lx (the great circle track) (by Prob. X.) from the pole S, and y will be the part of the track nearest to the south pole, or the *vertex*. Measure Sy (by Prob. XII.), and its complement will be the latitude of vertex.

By calculation :—

To find the two angles.

We have two sides and the included angle given.

a side	54°		
a side	50	included \angle	$2)110^\circ$
	<u>$2)104$</u>		<u>55</u> $\frac{1}{2}$ contained \angle
half sum	52		
half diff	$2)4$		<u>2</u>

To find the other angles (page 79).

as sine $\frac{1}{2}$ sum of sides 52°	. . .	co ar	0.103468
to sine $\frac{1}{4}$ diff of sides 2°	. . .		8.542819
so is cotang $\frac{1}{2}$ contained angles 55°	. . .		<u>9.845227</u>
to tang $\frac{1}{2}$ diff other angles	. . .		$8.491514 = 1^\circ 46' 30''$
as cos $\frac{1}{2}$ sum of 2 sides 52°	. . .	co ar	0.210658
to cos $\frac{1}{2}$ diff 2 sides 2°	. . .		9.999735
cotang $\frac{1}{2}$ contained angles 55°	. . .		<u>9.845227</u>
to tang $\frac{1}{2}$ sum of other angles	. . .		$10.055620 = 48^\circ 40' 00''$
		sum = $\angle x$	<u>$50 \ 26 \ 30$</u>
		diff = $\angle L$	$46 \ 53 \ 30$

Find Sy by rule of sines (78).


as rad	. . .		10.000000
to sine side 50°	. . .		9.884254
so is sine $\angle x$ $50^\circ 26' 30''$. . .		<u>9.887041</u>
to sine side Sy	. . .		$9.771295 = 36^\circ 12'$
			<u>90</u>

latitude of vertex "y" $\frac{53}{48}$

121. EXAMPLE 14.—To find an altitude of a celestial body.

Given latitude 30° N.
 „ time 11 A.M.
 „ declination 20° S.

Fig. 74.



By construction:—

Lay off the lat from R to P by scale of chords.

Draw the hour circle 11 A.M. = 15° by Prob. VIII. = $P \times S$.

Draw the parallel of declination $d e$ by Prob. VI. ; where these intersect will be x the altitude sought.

Through $Z x N$ draw a great circle (Prob. IV.), and $x y$ will be the measure of the altitude (Prob. XII.), and $P x Z$ is the triangle.

By calculation:—

With the co lat $Z P = 60^{\circ}$, the pole distance $x P = 110$, and the hour $\angle P = 15^{\circ}$, we have two sides and an included angle.

By rule page 79:—

a side	110°	
a side	60	
	2)170	sum
	85	$\frac{1}{2}$ sum
	2)50	the diff
	25	$\frac{1}{2}$ diff

2)15°	
7° 30'	half contained \angle

Draw a parallel of declination $dc = 20$ N. from E Q (Prob. VI.).

Through point of intersection x draw the great circle P x S (Prob. IV.), and P x R will be the triangle, and $x o$ the amplitude.

By calculation :—

P x will be a middle part (83), and the extremes are disjunct.

To find x R (89).

as $\cos 50^\circ$	co ar.	0.191933	
is to rad		10.	
so is cosine 70°		9.534052	
to cosine x R		9.725985	$= 57^\circ 51'$ azim
			90	
				the amplitude $x o = 32^\circ 9'$ N.

N.B.—The ordinary rule is, add the secant of the latitude to the sine of the declination, and the sine of the sum is the amplitude.

Thus,—sec 50°	0.191933
sine 20	9.534052
		9.725985 = $32^\circ 9'$

Remember the secant is the reciprocal of the cosine, and the cosine of $70^\circ =$ the sine of 20° ; hence the rule.

123. EXAMPLE 16.—To find the time of daybreak.

N.B.—Daybreak is the time at which the sun's centre is just 18° below the horizon of the place.


Given latitude	50°	N.
„ sun's declination	$9^\circ 43'$	S.

By construction :—

Lay off the lat 50° N. from R to P from scale of chords. Draw the parallel of declination dc by Prob. VI.

Draw the parallel tw of 18° below HR the horizon by Prob. VI.; where the parallels cut at x is the sun's place at daybreak.

Fig. 76.



Draw the great circle PzS (the hour circle), and ZxN the azimuth by Prob. IV., and ZxP will be the triangle, and the $\angle xPR$ the time from midnight.

By calculation:—

Here are three sides given, viz., the co lat 40° , the polar distance $99^\circ 43'$ and the side $Zx = 90^\circ + 18^\circ = 108^\circ$.

By rule, page 84. To find $\angle P$.

a side 108°	co arc sine	0.006275
99 43'	co arc sine	0.191933
40		
<u>2)247 43</u>		
123 52 $\frac{1}{2}$ sum		
99 43		
<u>24 9</u>	1st remainder sine	9.611858
123 52		
40		
<u>83 52</u>	2nd remainder sine	9.997507

$$\frac{1}{2} \angle ZPx = 53^\circ 15' = \frac{19.807573}{9.903786}$$

$$\angle ZPx = \frac{106 \ 30}{180}$$

$$\angle R.Px = \frac{73 \ 30}{4} \text{ in space by rule page}$$

$$60)294 \ 00$$

apparent time of daybreak = $4^h 54^m$ in time

N.B.—To be corrected by the equation of time for the day as given in almanacks, so as to get mean time.

If we use the triangle ZxP , we have a quadrantal triangle for $Zx = 90^\circ$.

By rule (page 69).

Find $\angle ZPx$.

as rad	10
is to cotang ZP	$38^\circ 33'$	10.098617
so is cotang Px	$100^\circ 15'$	9.257269
to cosine $\angle ZPx$	<u>9.355886</u> = $76^\circ 53'$ &c.

The preceding examples comprise all ordinary questions to which the attention of the navigator is likely to be called. When a student has read this little book he will be better able to comprehend works written upon the subject of spherics, in which the *theory* is explained. It is more than probable that a small volume may shortly follow the publication of this, adapted to those who, not content with an "*initiation*" into nautical astronomy, desire to become further acquainted with so beautiful, enticing, and useful a branch of study.




THE END.

LONDON

PRINTED BY SPOTTISWOODE AND CO.
NEW-STREET SQUARE

100



UNIVERSITY OF CALIFORNIA LIBRARY
BERKELEY

**THIS BOOK IS DUE ON THE LAST DATE
STAMPED BELOW**

Books not returned on time are subject to a fine of 50c per volume after the third day overdue, increasing to \$1.00 per volume after the sixth day. Books not in demand may be renewed if application is made before expiration of loan period.

JUL 25 1917

MAR 1 1918

FEB 27 1928

APR 08 1988

AUTO DISC MAR 11 1988

Saxby
Projection and calculation of the sphere

S3

Feb 20 1913

Lyon

FEB 22 1913

YB 04

FEB 22 1914

Jamleyson 24

MACAULAY.

MAR 25 1917

Robertson 7-5-174

8-4-17

HP

JUL 25 1917

MAR 1 1918

Wancheloge FEB 18 1918

Right Hon. Lo
Quarterly Magazine
and Historical Essays
ous Poems, &c. 2 v

FEB 27 1927

ACHS FEB 20 28

ession of James t
n, revised and correct

LAND from t
and II. 8vo. price 32

ontributed to t
Library Edition (t

RICAL ESSAY
Values for the Pock

RICAL ESSAY
tion, complete in 10
; call by HAYDAY, 3

RICAL ESSAY
ion, complete in 2 v

SAYS which ma

107275

LIBRARY

U.C. BERKELEY LIBRARIES



C006156362

PEECHES of the P
HIMSELF 8vo. 12s.

ORD MACAULA

III FORM in 1831 and 1832.

XI
AYS of ANCIENT ROME. By

MACAULAY. With illustrations, original and from the Antique, in
F.S.A., engraved on Wood by S. WILLIAMS. New Edition. Fcp. 10s. 6d.
marro; morocco, by HAYDAY, 42s.

XII
ORD MACAULAY'S LAYS of ANCIENT ROME, with IVR
and the ARMADA. 1832. 4s. 6d. cloth; morocco, by HAYDAY, 10s. 6d.

London: LONGMAN, GREEN, and CO. Paternoster Row.

